



US Army Corps
of Engineers



AQUATIC PLANT CONTROL
RESEARCH PROGRAM

MISCELLANEOUS PAPER A-69-3

ALFRED COFRANCESCO

REVIEW OF SENESCENCE AS AN IMPORTANT
FACTOR DETERMINING THE RELATIONSHIP
AMONG AQUATIC PLANTS, THEIR
EPIPHYTES, AND PATHOGENS

by

Eliska Rejmankova

Botany Department
University of California
Davis, California 95616



April 1989

Final Report

Approved For Public Release; Distribution Unlimited

Prepared for DEPARTMENT OF THE ARMY
US Army Corps of Engineers
Washington, DC 20314-1000

Monitored by Environmental Laboratory
US Army Engineer Waterways Experiment Station
PO Box 631, Vicksburg, Mississippi 39181-0631

Destroy this report when no longer needed. Do not return
it to the originator.

The findings in this report are not to be construed as an official
Department of the Army position unless so designated
by other authorized documents.

The contents of this report are not to be used for
advertising, publication, or promotional purposes.
Citation of trade names does not constitute an
official endorsement or approval of the use of
such commercial products.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.			
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S) Miscellaneous Paper A-89-3		
6a. NAME OF PERFORMING ORGANIZATION University of California	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION USAEWES Environmental Laboratory			
6c. ADDRESS (City, State, and ZIP Code) Davis, CA 95616		7b. ADDRESS (City, State, and ZIP Code) PO Box 631 Vicksburg, MS 39181-0631			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION US Army Corps of Engineers	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER			
8c. ADDRESS (City, State, and ZIP Code) Washington, DC 20314-1000		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Review of Senescence As an Important Factor Determining the Relationship Among Aquatic Plants, Their Epiphytes, and Pathogens					
12. PERSONAL AUTHOR(S) Rejmankova, Eliska					
13a. TYPE OF REPORT Final report	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) April 1989	15. PAGE COUNT 107		
16. SUPPLEMENTARY NOTATION Available from National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Aquatic plants Senescence		
			Epiphytes		
			Pathogens		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) This report is a review of literature on senescence and relationships between senescence and microorganisms in aquatic plants. The first part of the report summarizes the main facts regarding senescence as follows: a. Senescence is an important and not very well-understood stage in the life cycle of a plant. It is controlled by internal and external (environmental) factors. The precise mechanism of senescence induction is not known, but, in general, the initiation of senescence involves an imbalance in the relative levels of growth hormones. The change in hormone status can be caused by internal factors or environmental stimuli. b. Among the environmental stimuli involved in senescence initiation, the following were reported: light, day length, temperature, mineral nutrients, and pathogens. (Continued)					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL

19. ABSTRACT (Continued).

When pathogens attack plants, it is not always known whether hormonal imbalances are the cause or the result of infection.

- c. Senescence is accompanied by a change in enzyme activities. Peroxidases, glutamate dehydrogenase, endopeptidases, and some hydrolases have been reported to increase their activities during senescence. Dehydrogenase, ribulosebiphosphate carboxylase, glutamine synthetase, and glutamate synthetase generally decrease their activities during senescence.
- d. An extensive literature identifies cytokinins as the most generally effective senescence-retarding growth regulators. Abscisic acid (ABA) and ethylene, on the other hand, are known to be promoters of senescence. In many cases, cytokinins and ABA interact in a competitive manner.
- e. Decreases in chlorophyll content and protein amount are most often measured as senescence characteristics. Senescence may start long before changes in these parameters become apparent. No universal marker of senescence has been found to date.

The second part of the report describes epiphytic microorganisms on aquatic plants:

- a. Aquatic macrophytes often serve as a substratum for epiphytic microorganisms, namely algae, bacteria, and fungi. The amount and diversity of epiphytic microorganisms generally increase with the age of their host, mainly due to the increasing amount of organic matter excreted by the aging host plant.
- b. Microorganisms live mostly as commensals (saprophytes) on their hosts but may become parasitic as the hosts undergo stress or approach senescence.
- c. In submersed aquatic plants that usually have a very thin and reduced cuticle, bacteria do not need wounds for entry into plants. Bacteria often invade and degrade epidermal cell walls. Some are known to produce lytic enzymes. Some bacteria were reported as being able to produce antibiotics.
- d. Fungi are usually regarded as the main decomposers of dead organic matter in aquatic ecosystems. Many fungal species are known to produce enzymes capable of degrading cellulose, pectin, starch, etc. Besides enzymes, some fungi also produce toxins and antibiotics. The phytotoxicity of fungi has a potential use as bioherbicides. In this study, 150 fungal species were reported as being potential parasites on aquatic plants.
- e. Epiphytic algae seldom harm their host plants unless the colonization is so dense that it shades plants or, especially in combination with inorganic silt particles, it becomes so heavy that the host plants are damaged and dragged to the bottom.
- f. Few viruses were reported from aquatic plants. Viruses may offer an alternative for biological control, but at this point data are insufficient to even consider this possibility.
- g. Nematodes contribute to macrophyte degradation in certain cases by wounding the tissue and thus causing easy entry of infection. So far, at least 32 genera of nematodes that were parasitic on aquatic plants have been reported.

Two main points can be made from the literature search: (a) senescence is a process that can be induced by changing the environmental conditions, and (b) plants in the senescent stage are more susceptible to pathogens than young or mature but nonsenescent plants.

The report concludes with recommendations for future research.

PREFACE

This study was conducted as a part of the US Army Corps of Engineers Aquatic Plant Control Research Program (APCRP). Funds for this study were provided by the US Army Corps of Engineers (USACE), under Department of the Army Appropriation No. 96X3122 Construction General. The principal USACE Technical Monitor for the APCRP was Mr. E. Carl Brown.

The principal investigator for this work was Dr. Eliska Rejmankova, Botany Department, University of California, Davis, CA (UC Davis), who prepared this report. The study was conducted under the direct supervision of Dr. Lars Anderson, leader of the Aquatic Weed Research Group, US Department of Agriculture, UC Davis.

This research was monitored at the US Army Engineer Waterways Experiment Station (WES) by Dr. Alfred F. Cofrancesco, Jr., of the Environmental Laboratory (EL), Environmental Resources Division (ERD), Aquatic Habitat Group (AHG). The study was conducted under the general supervision of Dr. John Harrison, Chief, EL, and Dr. Conrad J. Kirby, Chief, ERD, and under the direct supervision of Mr. Edwin A. Theriot, Chief, AHG. Mr. J. Lewis Decell was Manager of the APCRP. This report was edited by Mr. Bobby Odom, while assigned to the Information Technology Laboratory under the Intergovernmental Personnel Act.

Commander and Director of WES was COL Dwayne G. Lee, EN. Technical Director was Dr. Robert W. Whalin.

This report should be cited as follows:

Rejmankova, Eliska. 1989. "Review of Senescence As an Important Factor Determining the Relationship Among Aquatic Plants, Their Epiphytes, and Pathogens," Miscellaneous Paper A-89-3, US Army Engineer Waterways Experiment Station, Vicksburg, MS.

CONTENTS

	<u>Page</u>
PREFACE.....	1
PART I: INTRODUCTION.....	3
Aquatic Plants and Their Control.....	3
History of Natural Macrophyte Decline.....	4
Management of Aquatic Weeds Based on Their Life Strategies.....	7
PART II: SENESCENCE.....	9
Background.....	9
Senescence and Disease.....	14
Senescence in Aquatic Plants.....	16
PART III: MICROORGANISMS ON AQUATIC PLANTS.....	28
Interactions Between Epiphytic Microorganisms and Their Macrophyte Hosts.....	28
Bacteria.....	31
Epiphytic Algae.....	35
Phytoplankton.....	39
Fungi.....	40
Viruses.....	43
Nematodes.....	46
PART IV: SENESCENCE AND PATHOGEN INTERACTION: ITS ADVANTAGES AND LIMITATIONS.....	50
PART V: CONCLUSIONS AND RECOMMENDATIONS.....	52
Conclusions.....	52
Recommendations.....	53
REFERENCES.....	55
BIBLIOGRAPHY.....	74
TABLES 1-5	
APPENDIX A: GLOSSARY.....	A1

REVIEW OF SENESCENCE AS AN IMPORTANT FACTOR DETERMINING
THE RELATIONSHIP AMONG AQUATIC PLANTS,
THEIR EPIPHYTES, AND PATHOGENS

PART I: INTRODUCTION

Aquatic Plants and Their Control

1. Even though some problems with aquatic plants existed in previous centuries, the main invasions and spread of nuisance aquatic plants in lakes and in navigation and irrigation canals worldwide have occurred in this century. The two primary reasons for this are (a) increasing water eutrophication and (b) the introduction of alien species into new areas. The time sequence of the most important aquatic plant invasions, together with the methods used for their control, was recently discussed by Anderson (1986).

2. For centuries people tried to control aquatic plants manually, perhaps with the exception of using herbivorous fish in China. With the development and registration of herbicides in the 1950's and 1960's, chemical control has replaced some manual methods. But the application of herbicides has not always been the solution. On the contrary, weed problems have sometimes even increased (Van Zon 1977). Some herbicides used for control of aquatic plants are selective, which means that other weeds usually replace those controlled. In stagnant waters, the decomposition of herbicide-killed plants leads to a further increase in eutrophication. The use of chemical compounds in water is requiring stricter regulations because of possible side effects and potential ground-water contamination. Unsatisfactory results with chemical aquatic plant control and increasing costs of herbicides have stimulated the search for new, alternative methods.

3. Many attempts have been made to regulate and restrict nuisance weed growth using either a selective biological agent (pathogenic bacteria and fungi, viruses, nematodes, insects, allelopathic plants) or herbivorous organisms. Several large-scale research programs on biological control of aquatic plants were started in the early seventies (Freeman et al. 1974, 1981; Bennett 1977). Many of them have been successful; for example, the fungus *Cercospora rodmanii* was used for control of waterhyacinth (Pennington and Theriot 1982),

and the alligatorweed flea beetle and alligatorweed stem borer were used as control agents for alligatorweed, *Alternanthera phytolaxeroides* (Cofrancesco 1984). Most recently, excellent control of *Salvinia molesta* in Australia was obtained with the introduction of the weevil *Cyrtobagous salviniae* (Room 1986). Many other projects that looked very promising in laboratory tests failed large-scale trials in the field. A new possible approach to biocontrol of aquatic plants seems to be the application of genetic engineering technology (Pennington 1985).

4. In the last few years, integrated management of aquatic weeds combining several control methods has been stressed. For example, the combination of insects (waterhyacinth weevils and waterhyacinth moths) with the parasitic fungus (*C. rodmani*) was used for controlling waterhyacinth in Louisiana (Sanders and Theriot 1985). These control agents have to be combined in such a way that they act synergistically. Insects often damage the leaves of aquatic plants and provide points of entry for pathogenic fungi or bacteria (Charudattan et al. 1978). Recently, Charudattan (1986) described integrated control of waterhyacinth using a pathogen, insects, and sublethal rates of herbicide. The US Environmental Protection Agency (EPA) sponsored several projects summarizing integrated pest control research and methodology (Tyndall et al. 1981; Aurand 1982, 1983; Trudeau 1982; Tyndall 1982).

History of Natural Macrophyte Decline

5. Dramatic changes have been observed in aquatic plant populations in the last several decades. New, introduced species have become established while others have disappeared. Increasingly disturbed aquatic ecosystems are more susceptible to sudden invasions. Introduced species often thrive at first, being free of natural parasites and foraging animals from their native locations. After a few years, however, a sudden outburst of growth is often followed by an almost equally sudden decline of most of the population; e.g., a rapid spread of *S. molesta* in Lake Kariba was followed by its degeneration (Loveless 1969), expansion of *Myriophyllum spicatum* in Chesapeake Bay in the late fifties to early sixties was followed by its decline in the mid seventies (Bayley et al. 1978; Orth and Moore 1984), and similar situations occurred in lakes in Wisconsin a few years later (Carpenter 1980b, Jones et al. 1983). A sudden decline of *Hydrilla verticillata* was described from Orange Lake, FL

(Haller 1983). Reasons for this reduction were not entirely known, but higher water level most probably played a major role.

6. Declines have been observed not only in populations of introduced aquatic macrophytes but in native species as well. In the 1930's an epidemic of the so-called "wasting disease" almost entirely destroyed the normally dense beds of marine eelgrass, *Zostera marina*, along the Atlantic coast of North America and Europe (Renn 1936, Rasmussen 1977). Motta (1978) described a considerable decline of *Potamogeton perfoliatus* and *Ruppia maritima* in Chesapeake Bay. *Myriophyllum spicatum* and *P. perfoliatus* stands are decreasing drastically in Neusiedlersee in Austria (Schiemer and Prosser 1976); Jupp and Spence (1977) described the decline in submersed macrophyte growth in Loch Leven. Increased occurrence of sudden macrophyte death was described in lakes in Norway (Rorslett 1985) as well as in a fjord in Denmark (Kiorboe 1980b). Marked regression has occurred in the growth of *Potamogeton pectinatus* in Swartvlei Lake in South Africa (Taylor 1983).

7. None of these declines has a simple cause, and none has been adequately explained. Declines always occur because of a combination of several interacting factors. Dozens of papers were published offering various explanations of drastic *Zostera* decline; the first and still most widely held view is that the mass destruction was the result of a disease caused by one or more parasitic organisms (the fungus *Ophiobolus halimus* or slime mold *Labyrinthula macrocystis*). It has never been proved that any of the organisms found in dead eelgrass can directly cause disease and subsequent death of the plant. After reviewing the literature on *Zostera* and the "wasting disease," Rasmussen (1977) suggested that temperature may play a decisive role in the destruction of the eelgrass. He also pointed out that once the plants are weakened by unfavorable temperatures (or any other environmental factor), then the so-far saprophytic microorganisms may become pathogenic and speed up the senescence and decay.

8. Decline in *M. spicatum* in Chesapeake Bay was believed to have been caused by viruses or virus-like particles (Bayley et al. 1968); two sets of symptoms were designated as "Northeast Disease" and "Lake Venice Disease." In later studies, Bayley (1970) was not able to identify the causal agent for any of them. Bean et al. (1973) were unable to produce symptoms of "Lake Venice Disease" either by direct inoculation of healthy plants with extracts from diseased plants or by growing diseased and healthy plants in the same

container. Only plants grown under low light intensity and inoculated with extracts from diseased tissue produced symptoms resembling those of "Lake Venice Disease." Plants growing under stress conditions (low light intensity) were apparently more susceptible to attack by microorganisms.

9. Carpenter (1980b) explained that no single factor can be responsible for the decline of *M. spicatum* in Lake Wingra, WI. He suggested that the most probable causes were increased turbidity, shading by epiphytes, and, possibly, increased susceptibility to parasites. Jones et al. (1983) presented a growth model of *M. spicatum* in Lake Wingra that predicted a substantial decline in macrophyte biomass due to phytoplankton increase. Available data do not show whether the phytoplankton increase preceded or followed the macrophyte decline.

10. Phillips et al. (1978) discussed the hypothesis that increasing phytoplankton biomass following eutrophication absorbs enough light to prevent net photosynthesis by aquatic macrophytes. These authors suggested, however, that the situation is more complicated. First, increasing epiphyte growth and increasing growth of a blanketing mass of filamentous algae will reduce the light intensity available to macrophytes so that a greater proportion of fixed energy is required for maintenance, and less is available for growth. Phillips et al. (1978) also considered the suppressive influence of organic secretions by the macrophytes on phytoplankton as shown, e.g., by Fitzgerald (1969). As their growth decreases, the influence of macrophytes on the phytoplankton diminishes, and increasing growth of phytoplankton will further reduce the light available to the macrophytes.

11. A resource allocation hypothesis for macrophytes decline was offered by Johnstone (1982). He suggested that a switch in resource allocation from vegetative growth to sexual reproduction, along with a limited ability of the mature stand to replace itself, ultimately causes the species to decline in abundance. If this hypothesis is correct, control methods that stimulate the vegetative growth of these plants (harvesting) may effectively prolong the aquatic weed problem.

12. Macrophyte development in a eutrophic lake in western New York (Nicholson 1981), influenced by repeated herbicide applications and mechanical cutting, showed that, indeed, species capable of vigorous regeneration (*M. spicatum*, *Ceratophyllum demersum*, *Elodea canadensis*) were not substantially affected by herbicides and in several years dominated over native *Potamogeton*.

Nicholson pointed out that species with characteristics like: (a) poor regeneration from detached pieces, (b) emphasized sexual reproduction, and (c) poor regrowth and healing when cut would be most vulnerable to mechanical cutting. When he compared the healing capacities of *M. spicatum* and native *Potamogeton*, he found that *M. spicatum* heals and elongates rapidly while *Potamogeton* often suffer local tissue deterioration. *Myriophyllum spicatum* is known to produce phenolic compounds (Planas et al. 1981) that are deterrent to herbivores and pathogens, which may be the reason for its local advantage over *Potamogeton*.

Management of Aquatic Weeds Based on Their Life Strategies

13. Throughout their life cycles, plants exhibit sensitivity to environmental factors and are capable of a range of responses to the environmental fluctuations. Different stages of plant development and different physiological processes may have different temperature and light optima and may require different photoperiods. Also, the sensitivity to infection by pathogenic fungi or bacteria is not the same throughout the plant's life cycle. Therefore, an understanding of the biology and physiology of problematic aquatic species is a necessary basis for their successful management.

14. Much information on the physiological ecology of aquatic plants already exists (Spence 1972; Barko and Smart 1981; Titus and Stone 1982; Barko et al. 1984, 1986; Agami and Waise 1985). Some of the papers refer to new management possibilities using photoinhibition of reproductive organ formation (Klaine and Ward 1984; Spencer and Anderson 1986a, b) or naturally occurring competitive plants (Yeo and Thurston 1984). Anderson (1986) outlines future approaches and principles in aquatic plant management based mainly on manipulation of external factors with the aim to disrupt the life cycle of a plant.

15. Senescence is an important and not very well-understood stage in the life cycle of a plant. With a few exceptions, not much is known about the factors that influence or trigger senescence in aquatic macrophytes. Reports indicate that senescent plants are probably most susceptible to infections by pathogens (Leopold 1961; Bean et al. 1973; Rejmankova 1979; Sharma 1985).

16. This report has been prepared with the aim of reviewing the available information on senescence of aquatic macrophytes and on microorganisms involved in the senescence process.

PART II: SENESCENCE

Background

17. Death of an organ or of the whole plant is always preceded by the process of senescence, which may be regarded as the final phase in development that leads to cellular breakdown and death (Wareing and Phillips 1985). More precisely, in leaves, it relates to the loss of structural integrity and photosynthetic competence of the chloroplasts. Senescence is also defined as the phenotypic manifestation of deleterious effects accumulating in old age, namely errors committed by the molecular genetic apparatus during protein synthesis (Leshem 1986). The term "senescence" has a somewhat confusing overlap in meaning with the term "aging." According to Leopold (1975), senescence refers to the deteriorative processes that are natural causes of death whereas aging refers to the wider array of processes of accruing maturity with the passage of time, without referring to death as a consequence. Of the steps in biological development, senescence is one of the least defined (Leopold 1961). Changes in growth rates and vigor and increases in susceptibility to environmental stress or pathogens are generally connected with senescence.

18. Before considering plant senescence further, it is necessary to distinguish between monocarpic species, which flower and fruit only once and then die, and polycarpic plants, which flower and fruit repeatedly. Monocarpic species include all annual and biennial plants and a certain number of perennial plants that grow vegetatively for a number of years and then suddenly flower, fruit, and die. Thus, in monocarpic species, death of the whole plant is closely connected with reproduction and is evidently genetically determined to occur at this stage of the life cycle. On the contrary, in polycarpic species, death of the whole plant is not normally associated with reproduction, and there is a great variation among the different individuals of a given species with respect to the length of the life span.

19. Another convenient distinction is between organ senescence and whole plant senescence. In most plants, each leaf has only a limited life span so that as the shoot continues to grow in height, the older leaves at the base tend to senesce and die. This pattern of senescence has been described as sequential senescence, and it must be distinguished from simultaneous or synchronous senescence (e.g., leaves of temperate deciduous trees).

20. Figure 1 illustrates the parts of a plant that can undergo senescence during its life. The submersed aquatic plant *M. spicatum* is shown in the picture.

21. Various internal factors of the plant are involved in the regulation of senescence (hormones, growth regulators), and a number of external factors may affect the rate of senescence, including light intensity and day length, mineral nutrition, disease, etc.

22. The precise mechanism of senescence induction is not known (Biswal and Biswal 1984). Factors influencing the initiation of senescence and the initiating events are described in detail by Thomas and Stoddart (1980), who




Figure 1. Parts of a submersed aquatic plant that can undergo senescence. Environmental and hormonal influences are indicated together with those parts of the plant that can exhibit correlative controls

emphasized the regulatory aspects of senescence and its characterization as a genetically directed process. These authors described the following factors:

- a. Genetic. Senescence is an integral part of plant development and, therefore, subject to direct genetic control. To what extent the control is directed from the genome of either the nucleus, chloroplast, or mitochondria has been a subject of many studies and discussions and is not clear yet (Osborne and Cheah 1982).
- b. Correlative. Senescence may be induced in a given organ (leaf) as a direct consequence of its position relative to other organs. It is a result of competition among different organs of a plant for light, space, translocated nutrients, and growth regulators.
- c. Environmental. Many experiments have been done in an attempt to assess the role of environmental factors in initiating or delaying senescence. Exposure to low light promotes senescence. Senescence is also influenced by day length. Exposure to adverse temperature in either the high or low range can bring about a senescencelike response, presumably as a consequence of structural or metabolic damage. Deficiency in mineral nutrition results often in chlorosis, necrosis, and leaf senescence. Often senescence appears to be induced in a direct response to a deficiency resulting in redistribution of the limiting nutrients from older tissues to developing structures. This is particularly a case of the mobile elements nitrogen, phosphorus, potassium, magnesium, sodium, and chlorine. The presence of pathogens within a leaf can result in acceleration of senescence.

23. Light is without any doubt one of the most important environmental factors, and its role in photosynthesis, plant development, and morphogenesis has been thoroughly studied. However, nothing definite is known about the mechanism of light action in controlling senescence although it has been known either to stimulate or retard this process. Biswal and Biswal (1984) reviewed the present knowledge on photocontrol of leaf senescence. Photostimulation of senescence is usually a question of high light intensities that may lead to photobleaching of chlorophylls. High light intensities not only accelerate chlorophyll breakdown but also lead to ultrastructural modifications and loss in soluble carbohydrates. The action of light in controlling senescence is normally attributed to its effect in delaying senescence. The action of light in retarding senescence has been attributed by many authors to its role in photosynthetic production of organic nutrients. Theories also suggest that phytochromes regulate leaf senescence. Another possibility is that light may induce changes in the level of phytohormones that influence senescence.

24. Senescencelike symptoms caused by nutrient deficiency (namely nitrogen) have been demonstrated in many plants. Jackson and Drew (1984) describe

the accelerated senescence in flooded plants. They stress nitrogen deficiency as a factor responsible for early senescence. Deficiency of nitrogen was suggested as being responsible for an early senescence of detached shoots of *Myriophyllum verticillatum* that were dependent on the nitrogen supply from nutrient-poor water as opposed to the rooted shoots that obtained enough nitrogen from sediments (Weber and Nooden 1974). High concentrations of CO₂ are known to increase the biomass production and accelerate senescence in terrestrial plants (Omar and Horvath 1983).

25. Present knowledge of the physiology and biochemistry of senescence is derived mainly from studies on leaves (both attached and detached leaves or leaf discs). Woolhouse and Jenkins (1983), however, warn against broad generalizations based on results of experiments that were done on relatively few plant species. The physiological responses, metabolic changes, or regulatory mechanisms involved in senescence will not be the same for plants adapted to totally different types of habitats (for aquatic macrophytes, a plant species growing in nutrient-poor oligotrophic lake with a high light penetration as opposed to a plant from nutrient-rich, eutrophic pond with a high turbidity).

26. The role enzymes play in leaf senescence was reviewed by Lauriere (1983). The main groups of enzymes studied were hydrolases, ribulosebiphosphate carboxylase (RuBPC) and the enzymes of the Calvin-Benson cycle, and a limited number of other enzymes, especially oxidoreductases. Increased cellular hydrolase activity has often been presented as one of the various manifestations of senescence, in accordance with the degradative nature of the process. Not all hydrolases, however, undergo an increase during the senescence, and again, there are differences among different plant species.

27. The enzyme with the most well-known control during senescence is RuBPC, which is responsible for net photosynthetic CO₂ fixation. The activity of RuBPC reaches its maximum at the time of completion of leaf expansion; then it declines to only about 5 percent of maximum activity at the time of advanced senescence (Batt and Woolhouse 1975). This level of enzyme activity represents the major loss of soluble protein during senescence (Friedrych and Huffaker 1980). It seems probable that the loss of RuBPC is the primary event responsible for the decrease of photosynthesis (Thomas et al. 1978).

28. Among the other enzymes, dehydrogenases have been studied most. A decrease in their overall activity is generally observed during the last stages of senescence. An increase in overall peroxidase activity during

senescence has often been reported. Current research is oriented toward the determination of the capacity of peroxidases to eliminate hydrogen peroxide, the production of which increases during senescence (Birecka et al. 1979) and which is toxic to plant cells. Nitrate and nitrite reductases have been little studied during senescence. A considerable increase in the activity of glutamate dehydrogenase has been shown by several authors. An increased activity of endopeptidases during senescence has been reported (Keist and Feller 1984).

29. Growth regulators such as cytokinins, auxins, gibberellins, abscisic acid (ABA), and ethylene are all known to play some role in senescence, but there is no firm knowledge on their modes of action. A major action of plant hormones in regulating senescence has been attributed to their activity in regulating the opening or closing of leaf stomata (Osborne and Cheah 1982).

30. Both cytokinins and gibberellins are known to delay senescence in many different plant species. This effect was described for excised leaves as well as for leaves attached to the plant (Smith and Grierson 1982). ABA, on the other hand, has been found to accelerate senescence in leaves of many species (Colquhoun and Hillman 1972). In most cases, cytokinins and ABA interact in a competitive manner.

31. There is contradictory evidence concerning the effect of ethylene on the rate of senescence. It has been shown that ethylene induces senescence in leaves of certain species and that it is also produced by senescing leaves of those species (Gepstein and Thimann 1981); however, ethylene may be without effect in another species. According to Osborne (1984), the amount of ethylene in some senescing aquatic plants increases, but whether or not the increased amount of ethylene causes the senescence in those plants is not known.

32. Manos and Goldthwaite (1975) demonstrated a seasonal senescence cycle in excised mature leaves with the senescence rate being highest in February and lowest in July. The intrinsic fluctuation in tissue senescence rate might be caused by hormonal or other factors. Leaves growing during the long days in summer could contain higher amounts of senescence inhibiting regulators or lower amounts of senescence accelerating ones. No change in sensitivity to hormones was applied externally.

33. Based on the available evidence, neither a decline in endogenous levels of any currently known senescence-retarding hormone nor an increase in

a senescence promoter seems to be the primary event in the induction of senescence; however, such changes may well be part of, or contribute to, a more complex induction system (Thomas and Stoddart 1980).

34. Although often observed as yellowing of chloroplasts, senescence is usually initiated long before any symptoms are visible. Friedrych and Huffaker (1980) found that chlorophyll loss did not begin until 14 days after the initial loss of RuBPC. In detached leaves, senescence symptoms become clearly observable usually about 48 hr after leaf detachment. The senescence of leaves often involves loss of chlorophyll, increase in carotenoids, decline in protein and RNA levels, enhanced proteolysis, hydrolysis of starch and other polysaccharides, and, in some cases, increased respiration. Many different methods have been employed in measuring the rate of senescence. None of them seems to be applicable to all plants.

35. Manos and Goldthwaite (1975) suggested that the reciprocal of the time to 50-percent chlorophyll breakdown is a good measure of the rate of the overall senescence process in leaf tissues. Colquhoun and Hillman (1972) used RNA and protein assays for assessing the effect of ABA on senescence in leaf discs of radishes.

36. It is generally accepted that senescence follows flowering and fruiting. There might be cases when some stress-inducing factors cause an early senescence that influences flower initiation. This was described for *Lemna gibba* grown under controlled conditions by Kandeler et al. (1974). Duckweeds do not flower frequently in field conditions; they reproduce mainly vegetatively. However, flowering was often observed in dense stands of duckweeds that were exposed to extremely high temperatures and that might also suffer a nutrient deficiency. Shortly after flowering, whole stands usually died (Rejmankova 1973). It could be that the plants started to senesce because of stress conditions and that flowering was triggered by a change in the metabolism of the senescing plant.

Senescence and Disease

37. Disease of higher plants is more than just an act of parasitism; rather, it is a deleterious shift in the biochemical and physiological processes required for normal cellular function, differentiation, and reproduction (Daly and Knoche 1976). No matter how well the undamaged parts of an

infected plant compensate for the local effects of a pathogen, eventually the physiological control systems fail to function properly, and senescence and decay take over. Three areas can be identified wherein the pathogen may modify natural leaf senescence (Habeshaw 1984): (a) transport and sink effects between leaf and pathogen and leaf and plant, (b) the supply of nutrients, and (c) the effect of growth hormones. Very little is known regarding the nature of hormonal changes in diseased plants. In many cases it is not known whether hormonal imbalances in diseased tissues are the cause or the result of infection. In other words, does the plant senesce because it was attacked by the pathogen, or was it attacked by the pathogen because it was already in a senescent stage and therefore more susceptible to pathogen attack?

38. The long-term effect of attack by a parasitic fungus resulting in acceleration of senescence was described by Butler and Simon (1971). The view that certain microorganisms can benefit from the natural senescence of their host is supported by results of Kirk (1984). There is a general belief that weak pathogens are able to advance senescence (Smedegaard-Petersen 1985). The experimental evidence for this phenomenon is rather conflicting. In many cases, senescence in infected plants is accelerated by the reduction in the rate of photosynthesis brought about by toxic substances produced by pathogen (Krishnamani and Lakshmanan 1976).

39. It is clear that although a number of interesting approaches to the problem of senescence in plants are being taken, it is too early to be able to present a single, overall hypothesis that will account for all the facts. In general it can be recognized that the initiation of senescence involves a change in the relative levels of growth hormones, and this change in hormonal status may be caused by either internal factors or environmental stimuli.

40. There are apparently two types of senescence. One is the senescence that proceeds as a natural part of the whole plant development and is primarily controlled by internal genetic and correlative factors. The other can be called enforced or induced senescence (Butler and Simon 1971) and reflects the changes caused by some unfavorable environmental factors (mineral deficiencies, insufficient light) or diseases. The main focus herein is on the later aspect of senescence and its possible manipulation. Why is manipulation of senescence wanted? The potential benefits to be gained are in terms of

plant productivity (crop plants) if the senescence is delayed. For aquatic weed problems, however, it would be beneficial to accelerate senescence (selectively if possible) and thus make the weeds more susceptible to disease.

Senescence in Aquatic Plants

41. Except for a series of papers by Jana and Choudhuri (1979, 1980, 1981, 1982a-g, 1984) and Kar and Choudhuri (1985, 1986) describing senescence in detail in both intact and detached leaves of submersed macrophytes *Vallisneria spiralis*, *Hydrilla verticillata*, and *P. pectinatus*, there has not been much more published on senescence in aquatic plants. Senescence is briefly mentioned in several production-ecological studies; some data exist on translocation of minerals before and during senescence. Osborne (1984) in her review on the role of ethylene in aquatic and semi-aquatic plants briefly discusses the role of ethylene in senescence.

Results from outdoor ecological studies

42. Natural senescence is an integral part of the whole-plant development. Therefore the complete growth cycle of a plant must be understood for an understanding of natural senescence. The growth pattern of submersed macrophytes usually corresponds to the generalized pattern given by Westlake (1965). The typical curve is sigmoid, but finally the biomass starts to decrease (see Figure 2). In temperate and subtropical climates there is generally a rapid biomass increase in the spring, with the maximum in the summer. The maximum biomass usually occurs at the time of flowering. Occasionally there may be two distinct biomass maxima during the growing season. After flowering and formation of seeds, submersed aquatic species with an annual mode of growth gradually senesce (Adams and Prentki 1982, Brock et al. 1985, Kulshreshtha 1982, Ram and Kapoor 1976, Rogers and Breen 1980). Senescence of the below-ground parts in species with extensive rhizomes and roots (e.g., *Nymphaeaceae*) is usually a much slower process than the senescence of the above ground parts (Brock et al. 1985).

43. Many submersed macrophytes in temperate climates appear to grow as annuals. Some species always develop overwintering organs (turions) before they die (e.g., *P. pectinatus*, *Potamogeton natans*, *Vallisneria americana*); others persist without special perennation organs (*E. canadensis*, *C. demersum*); and some vary according to the local conditions (*Myriophyllum*,




Figure 2. Hypothetical growth and metabolism curves for a plant with the annual mode of growth (—●— Biomass, ---▲--- current gross productivity, —■— current net productivity, —●— current respiration rate, —▼— death losses) (Westlake 1965)

H. verticillata). Carpenter (1980a), who studied nutrient release by three common macrophytes from Lake Wingra (*V. americana*, *M. spicatum*, and *P. pectinatus*), found that *P. pectinatus* senesces early in the season, while *V. americana* senesces late in the season. *Myriophyllum* shows more or less continuous leaching of organic material because there are always some senescent plants present in the population. The differentiated timing of the life cycle has been recognized as one of the strategies in sharing a habitat when niches are tightly packed. Another example of coexistence of several submersed macrophytes includes *Myriophyllum*, *Potamogeton*, and *C. demersum* (Crowder, Bristow, and King 1977).

44. Even if they are perennial, most of the submersed aquatic plants exhibit a typical annual mode of growth. They often produce special overwintering organs (turions, tubers, winter buds). The germination of these overwintering organs at the beginning of the growing season is usually very rapid. Nevertheless, these plants are still at a competitive disadvantage compared with those that overwinter in the form of the whole green shoots since the

latter are able to photosynthesize early in the spring. *Myriophyllum spicatum* is known to overwinter as a whole plant in many areas it invades. Its capacity for early growth at the beginning of the growing season is often cited as the reason for fast colonization and successful competition of native aquatic species.

45. Plants in northern temperate climates generally follow a unimodal pattern with a single growth maximum and relatively short periods of biological activity separated by a long period of inactivity due to severe winters. In warmer climates, a bimodal growth pattern prevails. This involves growth and maturation of two different age classes (overwintering shoots from the last year are usually responsible for the spring maximum, and new present-year shoots for the late summer maximum). Some plant species are apparently more flexible and are able to change their growth strategy in changing environmental conditions. Haag and Gorham (1977) and Haag (1979) give very instructive descriptions of different growth strategies of *Elodea*, *Myriophyllum*, and *Potamogeton* in Lake Wabamun in Alberta. They compare a part of the lake that freezes during the winter with a part receiving a thermal effluent. *Elodea* and *Myriophyllum* exhibit a rather flexible strategy compared with *Potamogeton*, which shows strong endogenous regulation of its growth. Two maxima were usually found for *M. spicatum* biomass in Lake Wingra during the years of its maximum spread. In 1977, when *M. spicatum* started to decline, there was only one maximum (Adams and Prentki 1982). Hannan and Dorris (1970) provide an interesting example of the development of submersed vegetation in natural conditions in a spring-fed river with almost constant water temperature as well as very uniform chemical composition. The macrophytes grew year-round with the gross photosynthesis dependent on light intensity.

46. There are not many studies on growth cycles and on the primary productivity of submersed macrophytes from tropical regions. Unni (1976) describes the production of several submersed species (*H. verticillata*, *C. demersum*, *Potamogeton crispus*, etc.) in a lake in central India. There are two distinct growth periods. The first one starts in January, and plants reach their maximum biomass in late March or April. The second growth period starts with the onset of rains in June, and the maximum biomass is reached by September. Following the maxima, some plants senesce and die, and some survive with a very slow growth till the following growth period starts.

47. Huebert and Gorham (1983), who studied the mineral nutrition of *P. pectinatus* under standardized environmental conditions, observed the striking seasonal periodicity in biomass production. They suggested that an internal seasonal regulation of the growth of *P. pectinatus* existed independent of obvious external environmental signals. Ram and Kapoor (1976) described this pattern in *C. demersum* grown from seeds: seed germination--differentiation--flowering--senescence, resulting in a sigmoid growth curve with the spread over a time period of 2 months.

Changes in mineral composition of plants during senescence

48. As the plant ages, it undergoes changes in its mineral composition. In perennials with underground storage organs, nutrients are translocated from below ground parts to the leaves at the beginning of the growing season. As the leaves senesce, the nutrients move in the opposite direction. Very often senescence appears to be induced in direct response to a deficiency, resulting in redistribution of the limiting nutrient from older tissue to developing structures such as young leaves and fruits. This is especially true for the mobile elements nitrogen, potassium, phosphorus, magnesium, sodium, and chlorine. Nutrients are not always translocated during senescence. In some species there is almost no translocation, and most nutrients are leached from disintegrating cells as senescence proceeds.

49. According to Prentki et al. (1978), 50 percent of the nutrients from senescent leaves of *Typha latifolia* are translocated into the below ground rhizomes. In eelgrass (*Z. marina*), both leaching and translocation are responsible for a decrease in total soluble organic matter, C and N in senescing leaves (Harrison and Mann 1975). As senescence started, there was only about one-third of nitrogen compared with the full-growth phase. Total organic matter dropped from 90 percent in young leaves to 28 percent in senescing leaves.

50. The changes in the contents of the main nutrients in *Phragmite* shoots during the growing season are summarized in Figure 3. The nitrogen content was highest in May, during the period of maximum growth. The sodium content reached a peak just before senescence and then fell sharply. The potassium content was highest at emergence and then fell steadily, with a sharp drop at senescence. Calcium levels increased to a peak in early July and then remained constant until senescence. The magnesium content remained




Figure 3. The amounts (mg/g^{-1}) of nitrogen (---○---), phosphorus (—▲—), sodium (---●---), and potassium (—●—) in shoots of *Phragmites communis* (Mason and Bryant 1975)

steady throughout. Many differences in the nutrient content of *Phragmites* and *Typha* can be detected (Boyd and Hess 1970, Kvet 1975, Dykyjova 1978), but the data shown in Figure 3 are reasonably representative for emergent macrophytes.

51. Typically, there is a seasonal decline in potassium content and then a sharper decrease during senescence. Besides emergent macrophytes, this pattern was also reported for submersed and floating species (Peverly 1985, Rogers and Breen 1980). High concentrations of potassium are required for the activity of many enzymes participating in intermediary metabolism and biosynthesis. Four main physiological and biochemical roles of potassium are recognized (Clarkson and Hanson 1980): (a) enzyme activation, (b) membrane transport process, (c) anion neutralization, and (d) osmotic potential. Barko and Smart (1981) hypothesized that lack of potassium might be a key element in triggering the senescence processes. Seasonal changes in chlorophyll content in several aquatic plant species followed a unimodal curve with a sharp decline corresponding to senescence (Ho 1979).

Induction of senescence in aquatic plants

52. Among environmental factors inducing senescence, temperature, light, day length, mineral nutrients, and heavy growth of epiphytes have most often been mentioned for aquatic plants, with sometimes conflicting evidence.

53. A decrease in water temperature was reported to induce senescence in *H. verticillata* and *Potamogeton berchtoldii*, resulting in abscission of winter buds and turions. However, in *P. crispus*, senescence was induced on *P. crispus* by increasing water temperature and day length (Sastroutomo 1982). Based on the results of experiments in greenhouse-heated tanks, Barko and Smart (1981) asserted that in *H. verticillata* and *Egeria densa*, higher temperatures accelerate senescence. Early senescence was observed in a dense stand of duckweeds, *L. gibba*, growing in a sheltered lagoon in a littoral of Nesyt Lake in Czechoslovakia (Rejmankova 1973). Water temperature was much higher in the lagoon than in the surrounding reedbelt where the duckweed grew more slowly but did not senesce until later in the growing season. Similar results were obtained for *L. gibba* growing in controlled conditions in the laboratory; life span was much shorter in conditions ensuring maximal growth as compared with the life span in less favorable conditions (Rejmankova 1979). Apparently, temperatures providing maximal growth also advance the onset of senescence. Comparison of growth of *M. spicatum* in a cooling reservoir at three locations with different water temperature showed differences in phenological advancement (Grace and Tilly 1976). The warmer the temperature, the more advanced phenology, which results in the earlier senescence.

54. Submersed aquatic plants are usually well adapted to low light intensities. Once the light intensity decreases under the compensation point, plants start showing symptoms of senescence. One of the reasons for decreasing light intensities is higher water turbidity caused by either anorganic silt particles or phytoplankton blooms (Spence 1972). A sudden increase in water turbidity occurred in Cayuga Lake in 1972 as a consequence of tropical storm Agnes (Oglesby and Vogel 1976). In some plant species, a temporary submergence in the dark does not induce senescence but only quiescence (i.e., a temporary suspension of growth). Quiescence can be immediately relieved by transfer from dark to light, as was shown experimentally by Quimby and Kay (1977) for alligatorweed.

55. High light intensities are also known to be responsible for early senescence, especially in plants previously adapted to low light, as shown under experimental conditions by Barko and Smart (1981).

56. In some plant species, the day length changes seem to be important in inducing senescence. Drew (1979) reported the summer reduction in chlorophyll content in a seagrass *Posidonia* as a sign of senescence induced by changes in the photoperiod. Spencer and Anderson (1986a) recorded a decrease in chlorophyll *a* and carotenoid content in *Potamogeton nodosus* after the switch from long-day to short-day conditions. *Potamogeton pectinatus* did not display similar changes (Spencer and Anderson 1986b).

57. There are reports on the shading effect of epiphytic algae causing the early decline of submersed macrophytes (Kiorboe 1980b, Adams and Prentki 1982). Kiorboe (1980b) compared the seasonal increase of the biomass of *M. spicatum*, *P. pectinatus*, and several other submersed species from two different localities in a fjord in Denmark. At one locality, all macrophytes except for the *Chara* were heavily overgrown with epiphytes. These plants had almost half as much biomass and started to senesce about 1 to 2 months sooner compared with those without epiphytes. The heavy growth of epiphytic microorganisms was reported as being responsible for early senescence in *P. crispus* (Rogers and Breen 1981).

58. Mulligan et al. (1976) fertilized small experimental ponds containing submersed macrophytes (*E. canadensis*, *M. spicatum*, *Chara*) with nitrogen and phosphorus. They found that except for *Elodea*, the other species were not able to grow in the heavily fertilized ponds but senesced and died. It is questionable whether the main reason for the macrophyte decline was too high a concentration of nutrients or a heavy phytoplankton bloom responsible for increased light attenuation.

59. Herbicides in sublethal concentrations may enhance senescence in aquatic plants. Even if most herbicide tests are oriented towards finding lethal doses, there are some papers describing sublethal concentrations (Fowler 1977), especially in combination with other types of control (Charudattan 1986).

Senescence and turion production

60. Some aquatic plants produce specialized buds that are important both in vegetative reproduction and in the survival of the species through periods of stress (e.g., winter in the north temperate regions). Several different

terms have been used for these buds such as winter-buds, turions, hibernacula, and resting buds. These vegetative propagules have lately received a lot of attention, namely in *Hydrilla* and *Myriophyllum*. The objective has been to define the conditions under which they are produced (Weber and Nooden 1974, 1976; Aiken and Walz 1979) and to find ways to prevent or limit their production (Klaine and Ward 1984, Spencer and Anderson 1986a). In *Hydrilla*, vegetative propagule production is stimulated by a shortened photoperiod and does not seem to be temperature dependent. Artificial illumination during the night prevents propagule formation. A short-day photoperiod enhanced vegetative propagule production in *P. pectinatus* and *P. nodosus* (Spencer and Anderson 1986b).

61. In *Myriophyllum verticillatum*, turion development is controlled by both photoperiod and temperature. Weber and Nooden (1974) noted that detached shoots form turions earlier than rooted plants. The effect of detachment could be a response to mineral nutrient deficiency in water. Rooted plants which absorb most phosphorus and nitrogen from the substrate that does not undergo any substantial seasonal changes in nutrient amount compared with water are not expected to have their turion formation induced by nutrient deficiency. After abscission of turions and after formation of tubers in *Hydrilla*, the vegetative shoots usually senesce and die. There is not always abscission of the propagules from the vegetative plant. *Myriophyllum exalbescens* turions differ from those of *M. verticillatum* in size and shape, but the main difference is that there is no abscission layer at the base of *M. exalbescens* turions. As the turions become heavier during the winter, they sink to the bottom with the senescent pieces of parent plants still attached (Aiken and Walz 1979). Not much is known about what happens with the vegetative shoots if they are prevented from forming turions.

Senescence studied on detached
leaves of submersed aquatic plants

62. Plant physiologists studying senescence are usually interested in a very narrow part of the whole process (e.g., the behavior of a single organelle or a specific enzyme). The test plant or leaf is chosen according to its suitability for that special type of research without consideration of reactions of the whole plant population in its natural environment.

63. One of the first senescence studies in which an aquatic plant was used as a test object was done by Yoshida (1961, 1970), who studied

chloroplast senescence in detached leaves of *Elodea densa*. Yoshida indicated that chloroplast senescence is closely associated with the nucleus. Makovetzki and Goldschmidt (1976) continued chloroplast senescence studies using detached leaves of *E. canadensis*. Their conclusions confirmed that chlorophyll senescence is under nuclear/cytoplasmic control, and chloroplastic proteins synthesis is not required for chloroplast senescence.

64. A series of papers has been published on the senescence phenomenon in aquatic macrophytes (*H. verticillata*, *V. spiralis*, and *P. pectinatus*) by Indian scientists Jana and Choudhuri (1980, 1981, 1982a-g, 1984) and Kar and Choudhuri (1985, 1986). The authors used mostly detached leaves for their experiments and occasionally compared their results with data on intact leaves. Even if one may doubt the direct applicability of results obtained on detached leaves to intact leaves/plants, these results still serve as an indication of possible changes that can be expected during senescence.

65. In their first paper on senescence Jana and Choudhuri (1980) compared the protein and chlorophyll content as well as enzyme activities in young, mature, and old leaves of *Vallisneria* and *Hydrilla*. Later, the same tests were repeated with *P. pectinatus* (Jana and Choudhuri 1982g). In all plants, the amount of chlorophyll and protein content was highest in mature leaves, lower in young leaves, and lowest in the old leaves. Protease and RNase showed increasing tendencies from young to old leaves; catalase activity decreased. Contrasting results were obtained with acid and alkaline pyrophosphatases; both decreased with leaf age in *Hydrilla* while they increased with age in *Vallisneria* and *Potamogeton*. In detached leaves, the enzyme behavior was not the same as in intact leaves. Kinetin in all concentrations tested was effective in delaying senescence in detached leaves. The concentration of 50 $\mu\text{g}/\text{ml}$ was the most effective. When comparing the metabolic decline due to aging in tested plants, the authors concluded that the decline in *Hydrilla* was slower than in *Vallisneria* and *Potamogeton*.

66. Jana and Choudhuri (1982a) also studied glycolate metabolism in the above-mentioned plant species. Glycolate (which is the substrate of photo-oxidation and the key enzyme of this process) and the glycolate oxidase were measured in different age groups of leaves and in detached leaves. Both glycolate content and enzyme activity were highest in mature leaves and decreased with aging. The activity of RuBP carboxylase also decreased with the onset of

senescence (Jana and Choudhuri 1982d), which is in agreement with results for terrestrial plant species (Friedrych and Huffaker 1980).

67. All three tested plants were found to produce an increasing amount of ethylene with aging (Jana and Choudhuri 1982e). Ethylene is known to be involved in the senescence of terrestrial plants. Production of ethylene was reduced by kinetin and enhanced by ABA in all three species.

68. In other experiments (Jana and Choudhuri 1981, 1982c, 1984), *Hydrilla* was more tolerant to heavy metals (mercury, cadmium, copper, and palladium) than were *Potamogeton* and *Vallisneria*. All tested heavy metals apparently hastened the onset of senescence, possibly, as the authors suggested, by impairing the general synthetic ability and membrane integrity of tissues of treated plants. Kar and Choudhuri (1985, 1986) measured senescence characteristics in detached leaves in light and darkness and following treatment with the polyamine spermine.

69. Strother (1984) measured the activities of acid and alkaline pyrophosphatases in senescing duckweeds, *Lemna minor*, grown in either complete or phosphorus deficient medium. Alkaline pyrophosphatase is supposed to be an enzyme most active in the conditions of highest metabolic activity. Acid pyrophosphatase has a role of phosphate scavenger and helps to maintain phosphate homeostasis. In *L. minor*, the activity of alkaline pyrophosphatase decreases during normal senescence; its decrease is accelerated by phosphorus deficiency. Acid phosphatase activity increases slightly during senescence. Under phosphorus deficiency, its increase is higher.

Senescence versus decomposition

70. Decomposition (decay) is a process immediately following senescence. It is defined as a degradation (usually microbial) of dead organic material. It is not always easy to distinguish between senescence and decomposition. Part of a leaf may be already dead and decomposing while the rest is still senescing. The distinction between senescing and decomposing plants or their parts is often somewhat arbitrary.

71. A continuum of certain symptoms is also typical for both processes. Organic matter is being released by senescing cells and continues to be released, more rapidly, after these cells die. Often the same microorganisms are present on senescing as on dead plants; their numbers and activities increase on dead plants. In most plant production, senescence and decomposition follow sequentially in any organ and may occur simultaneously during the

life span of a plant as older organs (particularly leaves) die, and new ones are produced (Rogers and Breen 1982). In a population, some plants will begin to senesce and decompose earlier than others. A good understanding of natural senescence and decomposition processes within the population of aquatic plants is, by all means, necessary for any attempts to manipulate these processes from outside.

72. Knowledge of the decomposition processes, especially of the rates of decomposition, is essential to an understanding of the role macrophytes play in aquatic ecosystems. Consequently, numerous studies describing decomposition of different aquatic plants have been published (Solski 1962; Nichols and Keeney 1973; Mason and Bryant 1975; Kistritz 1978; Karpati and Pomogyi 1979; Carpenter 1980a; Adams and Prentki 1982; Polunin 1984; Brock et al. 1985). The most commonly used method for studying decomposition is the so-called "litterbag technique," which involves the use of nylon bags filled with cut pieces of a plant. The bags are placed in water, and in successive intervals changes in dry weight and, occasionally, nutrients are measured. Unfortunately, in many litterbag experiments, healthy plants that had been previously killed (mostly by drying in the oven at 105° C or by freeze-drying) were used as an initial material for decomposition, which makes the results hardly comparable with natural field conditions.

73. Kudryavstev and Kudryavsteva (1980) compared dry weight loss during decomposition in *Potamogeton lucens* samples collected in different stages of its development. Samples originating from young healthy plants were found to have a much higher initial loss of dry weight than samples from already senescing plants. Similar results were obtained by Rogers and Breen (1982) with *P. crispus*. They compared decomposition, measured as the loss of weight and main nutrients, of naturally senescent plant material with that obtained when plants were artificially killed by drying. The authors found that the overall rate of weight loss from dried plants was slower than that of senescent plants even if the initial loss was more rapid for the dried plants. Howard-Williams and Davies (1978) measured the rates of dry matter and nutrient loss during decomposition of *P. pectinatus*. They used *P. pectinatus* shoots in a late senescent stage as an initial material for litterbag tests, and they even gradually shifted the bags down in the water column so as to simulate the conditions in a natural dying *P. pectinatus* stand.

74. The decomposition process can be generally described by an exponential curve (Best et al. 1982, Rejmankova 1979). The more detailed studies often point out that during the first days, the decomposition proceeds much faster than the predicted exponential curve. The initial deviation is most certainly caused by the physical effects of leaching. The decomposition rate is dependent on the type of plant material used for decomposition. The half times of decomposition vary between 200 to 400 days for emergent macrophytes (*Typha*, *Phragmites*) and are much shorter for floating and submersed macrophytes (*Lemna*, *Nymphaea*, *Myriophyllum*, *Potamogeton*), usually between 20 and 50 days. The rate of decomposition also depends on temperature; it was found to decrease exponentially with decreasing temperature (Rejmankova 1979).

75. Godshalk and Wetzel (1978) followed the fate of dissolved organic compounds (DOM) released from senescing and decomposing aquatic plants. The rate of metabolism of these compounds was dependent on temperature but mostly on the oxygen content of the water. In oxygen deficient waters, DOM accumulates, while in oxygen-rich waters, the metabolization to CO_2 is very efficient. A model to predict daily release of DOM and dissolved total phosphorus from three decomposing species (*M. spicatum*, *V. americana*, and *P. pectinatus*) in Lake Wingra was constructed by Carpenter (1980a). Sterry et al. (1985) described the major end products of microbial decomposition of two aquatic species (*C. demersum* and *Lemna pucicostata*) in a closed system after 42 days. Despite differences in their morphologies, habitats, and epiphytic flora, the decomposition in both species led to the same products: acetate, propanoate, butanoate, and hydrogen. The absence of methane indicated nonmethanogenic fermentation. The plants were allowed to decompose in an oxygenated system with the microorganisms originally present in their phyllospheres. The methanogenic bacteria that require a completely anoxic environment were unlikely to be present, which explains why fermentation did not proceed beyond the acid-forming phase.

PART III: MICROORGANISMS ON AQUATIC PLANTS

76. All microscopic plants and animals should be regarded as microorganisms, i.e., bacteria, fungi, unicellular algae, protozoa, and the smallest metazoa. Also, viruses are included in the group of microorganisms even if they are not independent organisms themselves. Microorganisms play a very important role in aquatic ecosystems. This review is mainly focused on those aquatic microorganisms that are growing on or attached to aquatic plants and are capable of damaging these plants either directly (pathogens) or indirectly (light limitation, physical damage).

77. There are several ways in which plant pathogens exert stress or influence on their host plants. They can produce enzymes that dissolve the cell wall (cellulose, pectinase activities) or produce surface-bound and extracellular polysaccharides, toxins, or phytohormones. Bacterial phyto-toxins were reviewed by Strobel (1977). The genetics of pathogenicity and selected aspects of molecular biology of microbial pathogens were reviewed by Panopoulos and Peet (1985). Only a few papers from the vast amount of literature on plant pathogens deal with aquatic plant pathogens (Zettler and Freeman 1972, Andrews 1980).

Interactions Between Epiphytic Microorganisms and Their Macrophyte Hosts

78. Many aquatic bacteria, fungi, or unicellular algae live within or on the surface of higher aquatic plants. The attached microorganisms are called "epiphytic" (epiphytes, epiphyton, periphyton--for details on terminology, see Appendix A (Glossary) and Sladeckova (1962)). They may use plants just as a physical support, or they may be commensals and derive all or some of their food from the food or metabolic products of their hosts. Generally, this does not involve any advantages or substantial disadvantages for their hosts. There are, however, other microorganisms that, as parasites, feed on the cells of their host, often producing toxic substances and thereby causing disease and often death. These are the pathogenic microorganisms. The distinction between commensalistic and parasitic microorganisms is not always very clear. Often fungi or bacteria may be present on a macrophyte for a long time without causing any symptoms in the plant and then may become parasitic when the

macrophyte is stressed when it is in a senescent stage. The term "opportunistic" is often used to describe this behavior. The term "saprophyte" is usually meant as an opposite to parasite, but because saprophyte refers strictly to organisms obtaining nutrients from dead organic material, one should use the term "commensals" for those epiphytic microorganisms feeding on dissolved organic matter excreted by their living hosts.

79. Microbial epiphytes have a variety of physiological relationships with their host plants. Besides being able to become pathogenic themselves, they can, on the other hand, reduce the inoculum potential of a pathogen either by simply occupying the area invaded by the pathogen or by producing antibiotics against the pathogen (Leben 1965, Blakeman 1971).

80. Most pathogens of aquatic plants contact their host passively with the aid of water. No chemotactic responses have been reported. To infect a plant, pathogens must penetrate a physical barrier (the cuticle and the cell wall of the plant). Most pathogenic bacteria and some pathogenic fungi enter their host passively through natural openings (wounds, stomata), but some have evolved sophisticated means for active penetration (Misaghi 1982).

81. There are many descriptive papers on epiphytic organisms, primarily algae, in aquatic ecosystems. They deal mainly with species composition (Jones and Mayer 1983), spatial and temporal distribution (Bowker and Denny 1980, Goldsborough and Robinson 1985, Stowe 1982, and others), and biomass production (Bulthuis and Woelkerling 1983, Howard-Williams and Liport 1980, Penhale 1977, and others) of periphyton. Many studies that were done on periphyton did not even consider a host plant as an important part of a microorganism-macrophyte complex and used artificial substrata for evaluation of periphyton growth (Cattaneo and Kalff 1978).

82. Much less information is available on the physiology, nutrition, energetics, or pathogenicity of epiphytes in an aquatic ecosystem and on actual interactions between macrophyte and attached microorganisms. Some macrophytes do not have periphyton because of the release of compounds toxic to algae, (e.g., as described by Anthoni et al. (1980)). More commonly, however, dense and productive periphyton communities of algae and bacteria, often embedded in a carbonate muco-organic complex, develop on most macrophytes. Shading and nutrient depletion at the leaf surface are the major effects on the macrophytes. Heterotrophic microorganisms and algae can use dissolved organic carbon released by the macrophyte. The metabolic activity of the

heterotrophs, in turn, promotes the growth of algae and, probably, of the macrophyte by increasing the rate of mineral cycling (Sondergaard 1983). Pip and Robinson (1982) stressed the importance of the physical arrangement of epiphytic organisms and their host relative to each other.

83. Figure 4 shows interactions between macrophytes and their epiphytes. A very thorough study on primary productivity, chemoorganotrophy, and nutritional interactions of epiphytic algae and bacteria on macrophytes in the littoral region of a small temperate lake was done by Allen (1971). Allen's results confirm that DOM released extracellularly by macrophytes is rapidly incorporated into the epiphytic complex. Based on a series of experiments with axenic bacterial, algal, and macrophyte (*Najas flexilis*) cultures, Allen showed that organic materials released by *Najas* were not wholly suitable as carbon and energy substrate for algae without prior bacterial degradation. Allen also described how the colonization of a macrophyte proceeds. During initial colonization, bacteria adhere and probably form a monocellular layer. This is subsequently infiltrated with calcium carbonate precipitated by the macrophyte and algal forms, particularly in calcareous waters. As colonization intensifies (late spring and early summer), a dissolved organic carbon "pool" is probably established in the matrix of deposited carbonates. Its




Figure 4. Generalized pattern of interactions in a macrophyte-epiphytic microflora complex

sources are: (a) extracellular release from macrophytes, (b) active excretion from attached algae and bacteria, (c) decomposition products following autolysis of epiphytes, and (d) dissolved carbon compounds present in littoral water. Wetzel and Allen (1970) suggested a metabolic model of macrophyte-periphyton interactions. Its validity was confirmed by Allanson (1973), who studied the periphyton structure on *Potamogeton natans* and *Chara*. The relationships between epiphytic microfloral populations and the substrata upon which they grow were reviewed by Wetzel (1983). He admits that much is still unknown, and he stresses that in situ metabolism at the microlevel must be analyzed as well as quantification of the metabolic pathways by direct experimental methods.

84. A microcosm consisting of several types of microorganisms was described in internodal areas of *Spartina alterniflora* (Gessner et al. 1972). At first, fungal mycelia of *Sphaerulina pedicellata* with characteristic hyphopodia develop on the surface of growing *Spartina*. Hyphopodia serve for attachment, penetration, and absorption. As the fungal biomass increases, additional microorganisms appear, namely bacteria, such as *Metallogenium*, *Hyphomicrobium*, and *Mycoplasma*. These bacteria are known to grow only in the presence of fungi. The occurrence of the nematodes of *Oncholaimus* among the fungal mycelium as well as the presence of acarid mites is typical. During the senescence of *Spartina*, the fungal growth accelerates.

Bacteria

85. As was the case for other epiphytes, epiphytic bacteria have been the subject of mainly descriptive studies. Bacteria have been enumerated and occasionally their biomass estimated, but not much is known about their functions and interactions with host plants. They are regarded as initial colonizers of aquatic plants using organic matter exuded from macrophytes or released by aphid or other insect action from subcuticular tissues (Lee 1980). However, the evidence of the sequence of microfloral succession on aquatic plant material has been controversial (Robb et al. 1979). Many earlier studies regarded fungi as primary colonizers on decaying plants (Gessner et al. 1972).

86. The density of epiphytic bacterial populations in aquatic ecosystems depends on various factors. It tends to increase as a result of bacterial attachment and growth whereas expansion of plant tissue, detachment of bacteria, predation, and death can cause a decrease. Hossell and Baker (1979a, b) estimated average bacterial numbers on the lower surface of *Lemna minor* leaves and on *Rorippa* leaves. They found that bacterial density increased significantly with leaf age. Patience et al. (1983), who studied the decomposition of *L. paucicostata*, provided evidence that the microorganisms involved in decomposition are mainly rod-shaped or coccoid bacteria which are present in high densities on the lower surfaces of living *Lemna* leaves. Spatial distribution of epiphytic bacteria on seven species of aquatic plants was described by Baker and Orr (1986). The authors confirmed that the older leaves have much higher bacterial density. Two investigated species (*Nasturtium officinale* and *Veronica beccabunga*) were found to have significantly larger populations on the abaxial surfaces of submerged leaves than on the adaxial surfaces. The difference may be due to the unilateral release of hydroxyl ions during bicarbonate utilization.

87. Rhizosphere bacterioflora of *L. minor* was described by Coler and Gunner (1969); that of *Myriophyllum heterophyllum* by Blotnick et al. (1980). Both papers confirmed that microbial activity is enhanced by exudation of organic substances from aquatic plants. Density and production estimations of bacteria on several macrophyte species were presented by Kudryavstev (1984) and Kudryavstev and Yeshov (1980). They found increasing numbers of cellulosytic bacteria from the spring to the fall. Among the macrophyte studied, pondweed (*Potamogeton*) had the most abundant bacterial epiphytes.

88. Rogers and Breen (1981) studied the epiphyton colonization of the leaves of *P. crispus* and tried to establish its effect on the host plant. They did not find any differences in colonization of abaxial and adaxial surfaces of the leaves. The youngest leaves were devoid of epiphyton. The young leaves were colonized by groups of short stout bacteria that appeared to be attached to the leaf surface by weblike structures. With increasing leaf age, the epiphyton became denser and included a greater variety of species. Besides short stout bacteria, some cyanobacteria were present as well as some bacilli and several diatoms species. The oldest leaves did not show any changes in species diversity, but the epiphyton increased in density. The electron microscopy revealed that bacterial attachment did not damage the

surface of young leaves. There were, however, extensive inward swelling and disorganization of the epidermal walls, characteristic of a reaction to invasion by pathogens. In older leaves, the swelling was also present in mesophyll cells while bacteria had invaded and degraded the epidermal cell wall. The bacterial invasion was concomitant with signs of senescence.

89. In a later study, Rogers and Breen (1983) included grazer interaction in their studies. The host plant (*P. crispus*) was resistant to consumption by snails until it was invaded by epiphytic necrotrophic bacteria, which made the host tissue more palatable for the grazer. Grazing of epiphytes by the snails, however, reduced the rate of development of the bacteria and probably extended the life of the plants. A succession of microflora on developing leaves of *P. pectinatus* was examined by Howard-Williams and Davies (1978). They found that prosthecae bacteria developed after 6 weeks of leaf growth. These bacteria were associated with cuticular erosion and epidermal pitting. Several weeks later the senescing leaves showed extensive cuticular peeling and epidermal cell wall rupture. Details of the forms of bacterial attachment to leaf surfaces of *P. pectinatus* were described by Robb et al. (1979). The authors assayed individual isolate cultures for cellulase activity and found that six out of eight isolates produced extracellular cellulase.

90. Bacterial attachment proceeds in the following stages: (a) a reversible ionic attraction, (b) secretion of an adhesive acidic polysaccharide that develops either into a fibrous web or a network of fimbriae, and (c) the formation of clumps. These steps seem to be necessary morphological adaptations in an aquatic environment; they allow for concentration of exoenzyme activity on a food substrate. Adhesion-enhancing materials and their specificity for bacteria-host interactions were described by Imam et al. (1984). It is generally believed that the entry of bacteria into plants requires wounds because epidermal cell walls are protected by cuticle (Kolattukudy 1985). The above examples show that this is not always the case in aquatic plants, presumably because the cuticle in submerged macrophytes is extremely thin and reduced (Sculthorpe 1967).

91. Many papers describe the epiphytic microflora on seagrasses. Kirchner (1984) measured dissolved organic carbon excretion rates by *Z. marina* leaves. It equaled 2 percent of total carbon fixation, and the epiphytic bacterial population was supported almost entirely by carbon obtained directly

from *Zostera* photosynthesis. Bacterial colonization was much higher at the tips of the leaves (*Zostera* leaves grow at the base, which means that leaf tips are older). The doubling time for bacterial population was 8 days. According to Waite and Mitchell (1976), the bacteria living on the surface of *Ulva lactuca* leaves act as typical opportunists. Under normal conditions they provide essential growth factors for their host, but they are capable of degrading *Ulva*. The degradation process is presumably triggered when *Ulva* gets into stress conditions. Lemos et al. (1985) isolated 224 bacterial strains from marine green and brown algae; 38 of these strains, all from the *Pseudomonas-Alteromonas* group, displayed antibiotic activity.

92. Waite and Kurucz (1977) isolated bacteria from decaying *Hydrilla* plants and tested them for the ability to produce cellulolytic enzymes. Those isolates able to degrade cellulose were used as inoculum for the degradation of *Hydrilla*. The plant material was autoclaved prior to inoculation, and the kinetics of plant material degradation was measured.

93. In recent years, attempts have been made to use epiphytic microorganisms to produce lytic enzymes for potential control of nuisance aquatic weeds. Gunner (1983) studied cellulolytic properties of enzymes produced by microorganisms occurring naturally on the surface of watermilfoil species. The principal microflora was bacterial. The author tested individual isolates as well as their mixtures and concluded that a mixture of microbial isolates was more effective in attacking the plant than the isolates individually. Pennington (1985) conducted a search for microorganisms among endogenous microflora of hydrilla that would be able to increase their production of lytic enzymes and hence be able to attack plant tissues. She did not find any cellulase bacteria and only one pectinase-producing bacteria isolate.

94. Ramsay and Fry (1976) and Fry and Ramsay (1977) studied the response of epiphytic bacteria from the treatment of *E. canadensis* and *Chara vulgaris* with the herbicide paraquat. The number of epiphytes on *Chara* that was resistant to paraquat did not change after the treatment. Epiphytes on *Elodea* that died after 15 days following the treatment had all increased in number in the following 10 days; apparently their growth was stimulated by the soluble organic matter released from dying *Elodea*. The difference in the number of epiphytes before and after the treatment was greater for *Elodea* plants that were healthy prior to the treatment compared with those that were initially moribund.

95. Table 1 gives several examples of bacteria described from aquatic plants. Many so-called "low-grade" phytopathogens are among bacteria. These phytopathogens usually attack plants that are under stress. Several *Pseudomonas* species belong to these low-grade phytopathogens. Several examples of *Pseudomonas* isolated from different aquatic plants are shown in the table.

Epiphytic Algae

96. Numerous papers have been published on algal epiphyton, but most of them have paid little attention to the substrate. On the other hand, many studies of aquatic plants note the presence of epiphyton without going into any details. Complex studies involving both epiphytic algae and their host macrophyte are scarce, especially those that evaluate what changes occur in a macrophyte as a consequence of periphyton colonization.

97. Periphyton studies are often complicated by methodological problems arising when periphyton has to be removed from its substratum. A review of methods of periphyton measurements was prepared by Weitzel (1979). An up-to-date evaluation of suitable methods and techniques has been presented by Delbecque (1985), who commented on the many different techniques used from simple shaking or scraping the host plant free of its periphyton to complicated bleaching techniques or enzymatic removal of particularly prostrate species. Apparently each approach requires a special technique, and one solution for all problems cannot be given. Carter (1982) described a technique for direct microscopic observation of periphyton on macrophytes. Another controversial issue is how to express the amount of periphyton. Many authors try to relate the periphyton biomass to the unit of leaf surface area of the host. In species like *Myriophyllum* or some *Potamogeton*, the leaf morphology is so complicated that determinations of average surface areas can usually be only a rough estimate. Therefore, expressing the periphyton amounts in terms of dry weight of macrophyte hosts seems to be less erroneous (Gough and Woelkerling 1976).

98. A common practice in studies of epiphytic communities is the use of artificial substrata. Seasonal changes in the epiphyton community on natural macrophytes (*Potamogeton richardsonii*) and on a similar plastic plant were compared by Cattaneo and Kalf (1978). The authors concluded that there were no significant differences in community composition in the early season; the

differences that developed subsequently appeared to be principally a function of the amount of CaCO_3 incrustation. In their further study, Cattaneo and Kalff (1980) evaluated the relative contribution of macrophytes and their epiphytes to the total production. *Myriophyllum spicatum*, *P. richardsonii*, and *V. americana* were compared and, again, artificial plants were used as well as natural ones. *Myriophyllum*, a species with the most highly dissected leaves, supported greater epiphyte biomass than the other plants. Cattaneo and Kalff (1978, 1980), as well as Fontaine and Nigh (1983), were convinced that over most of the growing season, macrophyte hosts serve as a neutral substratum for their respective epiphytes. Only during periods of macrophyte senescence caused by metabolite leakage may macrophytes become relatively more important in structuring epiphyte communities.

99. In strong opposition against the "neutral substrate" theory are O'Neil, Morin, and Kimball (1983) and Wetzel (1983), who argue that by placement of the artificial substrata used in the above-mentioned studies in situ in the macrophyte stands, periphyton colonizing the artificial substrata was exposed to macrophyte-induced changes in the surrounding aquatic environment.

100. Many algal epiphyton studies deal exclusively with diatoms that often are a major contributor to an epiphytic community. Stowe (1982) described the development of the diatom community on *S. alterniflora*. The maximum diatom numbers occurred during the period of *Spartina* standing crop decline. Leachates released from dying grasses apparently provided nutrient stimulation for diatom growth. Seasonal succession of diatom epiphyton on *L. minor* was studied by Goldsborough and Robinson (1985). Maximum diatom biomass was recorded in early spring while the diversity reached its maximum during the decline of duckweed biomass. The extensive bloom of diatoms on decomposing *M. spicatum* under the ice in Lake Wingra was reported by Boylen and Brock (1974). A distribution of epiphytic diatoms on the lower side of leaves of *Nuphar lutea* and *Nymphaoides alba* was described by Delbecque (1983). No differences in diatom colonization were found between the two nymphaeid species.

101. Macrophyte-mediated changes in the abundance of algal epiphyton were described by O'Neil, Morin, and Kimball (1983) for epiphyton on *M. heterophyllum*. The dense growth of *Myriophyllum* influenced physical and chemical conditions in the surrounding water column. Periphyton was dominated by diatoms in the early summer when the macrophyte occupied only the lower part of the water column. By midsummer the entire water column was occupied by

M. heterophyllum whose metabolic and photosynthetic activity and dense foliage created marked vertical gradients in physicochemical conditions. Periphyton species composition shifted to blue-green and filamentous algae on the apex and midstem. On the lower stem, diatoms consistently dominated the periphyton. Spring diatom maximum followed by a dominance of blue-green and filamentous algae in summer was also reported by Marvan et al. (1978).

102. Numerous papers describe algal epiphytes on brackish and marine submersed macrophytes. As these macrophytes are important as the primary producers in shallow tropical and temperate regions, they have been a subject of many production ecological studies (Harlin 1973, 1975; Ballantine and Almodovar 1977; Penhale 1977; Congdon and McComb 1979; Jacobs and Noten 1980; Willcocks 1982; Bulthuis and Woelkerling 1983; Moriarty et al. 1985; and others). Most of these authors agree that epiphyte colonization is heaviest on the oldest parts of leaves. Howard-Williams and Liport (1980) found very high epiphyte biomass on leaves of *Zostera*, often exceeding the biomass of the leaf itself. The role of epiphytes in the primary productivity of *Zostera muelleri* in Westport Bay, Australia, was described by Clough and Attiwill (1980); the epiphytes themselves contribute to the biomass production, but at the same time, they shade their host plant, thus apparently lowering its photosynthesis. Heijs (1985a) measured the biomass contribution by epiphytes to be as high as 35 to 44 percent of the total aboveground production. According to Verhoeven (1980), the shading of epiphytes does not really suppress the macrophyte photosynthesis substantially because the most dense epiphyton develops on the older leaves that are not so much photosynthetically active anymore. Johnstone (1979) found that epiphyte colonization restricts the photosynthetically active life of a leaf segment to less than 25 days. Sometimes grazing can contribute to the regulation of epiphyton growth and to maintaining the viability of seagrasses (Van Montfrans et al. 1982).

103. The importance of epiphytic algae in nutrient cycling (especially phosphorus and nitrogen) was described by Allen (1971) and Mickle and Wetzel (1978). Riber et al. (1983) pointed out that a considerable proportion of the phosphorous pool of a lake passes through the algal periphyton during the year. Many authors also reported on the excretion of organic carbon and its subsequent utilization by epiphytes (Wetzel and Penhale 1979).

104. Physical and chemical parameters regulating growth of algal epiphytic communities were discussed by Sand-Jensen (1983). The author

stressed that while the physicochemical parameters above the epiphytic community can be measured reasonably well, the conditions within the community are often quite different. This is true particularly for thick and dense epiphyton communities in standing waters. Nevertheless, simple measurements can be done relating the physical (light, temperature) and chemical (inorganic nutrients) factors to the composition and biomass of epiphytic algae.

105. The epiphytes are capable of reducing the photosynthetic rate of their host by reducing the light intensity (Penhale 1977). Sand-Jensen (1977) measured the light intensity reduction of about 45 percent by a thick crust of epiphytes, mainly diatoms. In a further study, he measured that in a lake with a high nutrient supply, epiphytes were responsible for 86 percent of light attenuation. Selective light attenuation by the periphyton complex was characterized by Losee and Wetzel (1983). Debusk and Ryther (1981) observed lowering in the productivity in unharvested cultures of *H. verticillata*. They explained it as a result of severe light limitation on a dense *Hydrilla* mat. They noted an extensive epiphyton growth that was also apparently responsible for restricted growth.

106. Several examples have been described where epiphyton was so heavy that it actually dragged down and submerged aerial shoots of submersed macrophyte species (Schiemer and Prosser 1976; Nuttall 1985; Nuttall et al. 1985). Organic and inorganic silt particles that stick to filamentous or meshlike algae may contribute substantially to the weight of the epiphyton complex. Menzie (1979) found as much as 3.8 g of particulate material attached per gram of *M. spicatum*.

107. The effect of increased water temperature resulting from thermal effluent on the growth of epiphytic algae was described by Hickman and Klarer (1974). An extended growing season was responsible for early epiphyton development, but temperature alone was not the reason for all the difference. Day length and light intensity were also responsible.

108. Hodgson and Carter (1982) estimated the effect of *Hydrilla* herbicide treatment on a periphyton community. They used glass slide periphyton samplers instead of assessing periphyton directly on *Hydrilla* plants. Nevertheless, the growth of periphyton was higher following the treatment compared with the untreated control. Anderson (1981) described the changes in species diversity of periphyton in a small macrophyte pond following a treatment by

hexazinone. The species diversity of periphyton was dramatically decreased after the treatment.

Phytoplankton

109. Numerous papers refer to the relationship between macrophytes and surrounding phytoplankton with often contradictory conclusions. Hasler and Jones (1949) were among the first limnologists to study this relationship in outdoor experiments. According to their results, a dense stand of submersed macrophyte *E. canadensis* significantly inhibited the growth of phytoplankton. Hogestu et al. (1960) also described an antagonistic relationship between rooted aquatic macrophytes (*Hydrilla*, *Vallisneria*, *Nymphoides*) and phytoplankton. The inhibition of phytoplankton growth was explained by shading effect and competition for nutrients plus some other unknown factors. Fitzgerald (1969) concluded that even if the main reason for the antagonistic relationship is based upon nutritional factors (nitrogen-limited cultures of macrophytes will remain relatively free of algae), there are indications of antibiotic factors as well. Planas et al. (1981) have shown that phenolic compounds extracted from *M. spicatum* can inhibit the growth of algae in uni-algal cultures and in natural assemblages in the laboratory.

110. Besides papers describing the limiting effects of macrophytes on phytoplankton, others report the inhibition of macrophyte growth by the abundance of phytoplankton. Jones et al. (1983) suggested that a major increase in phytoplankton abundance that resulted in a marked decrease of light penetration might be, at least partly, responsible for the decline of *M. spicatum* in Lake Wingra, WI. The growth of the submersed macrophyte *Zannichelia peltata* can be significantly inhibited by an algal filtrate (Van Vierssen and Prins 1985). Algal culture consisted mainly of *Anabaena*, and it was concluded that this alga released an unspecified allelopathic substance. Sharma (1985) found that the growth of *Eichhornia crassipes* was inhibited when grown in water containing algae such as *Scenedesmus bijugatis* and *Chlorella pyranoidosa*. Waterhyacinth plants ultimately died after 90 to 100 days of growth with these algae. The death of the plants was suspected to be due to allelopathic interactions between algae and waterhyacinths. The fungus *Alternaria eichhorniae* was isolated from the dying plants. The waterhyacinth may have become

susceptible to the fungus because it was weakened by the allelopathic effect of algae.

111. Results of experiments on the interaction between *M. spicatum* and phytoplankton indicate that *M. spicatum*, especially at the early stages of its growth, stimulated the growth of the phytoplankton (Godmaire and Planas 1983). One possibility is that extracellular organic matter released by *Myriophyllum* could be used directly or could influence indirectly the growth of phytoplankton. In their recent study, Godmaire and Planas (1986) confirmed their previous results. Multivariate analysis results showed that differences in phytoplankton productivity among sites with and without *Myriophyllum* were unrelated to nutrients and light. A factor not measured in their study must have been responsible for stimulation of phytoplankton growth. Landers (1982) studied the effect of naturally senescing aquatic macrophytes (e.g., *M. spicatum*) on nutrient chemistry and chlorophyll *a* of surrounding waters. He observed a substantial increase of phytoplankton because of organic substances released by senescing *Myriophyllum*.

Fungi

112. Among the main groups of fungi (*Myxomycetes*, *Ascomycetes*, *Basidiomycetes*, and Fungi imperfecti), there are some aquatic species. As zoospore producing organisms, fungi are well adapted to infect submersed aquatic plants. Until the early seventies, there were only occasional reports on fungal infection of aquatic plants, but no extensive, systematic research had been done. The increasing interest in biological control of nuisance aquatic weeds stimulated a research program on biological control of aquatic weeds at the Commonwealth Institute of Biological Control in Bangalore, India (Zettler and Freeman 1972). In 1970 a project on the evaluation and subsequent use of plant pathogens as biocontrol agents for aquatic weeds was begun at the University of Florida, Gainesville (Freeman 1977). Both endemic and exotic plant pathogens (mostly fungi) were collected and tested (Freeman et al. 1974, 1981, 1982; Freeman 1979). The Aquatic Plant Control Research Program (APCRP) has been coordinated by the US Army Engineer Waterways Experiment Station (WES), Vicksburg, MS.

113. Table 2 presents a list of potentially pathogenic fungi that were reported from aquatic plants. In Table 3 the same fungi are arranged

according to their macrophyte hosts. Besides freshwater species, *Spartina* and seagrasses from brackish waters (*Zostera*, *Ruppia*) were included. Only several selected examples of fungal microflora on *Spartina* are presented. Gessner and Kohlmayer (1976) list about 100 fungus species that have been reported from *Spartina*. As this plant is not a central point of interest, these fungi are not included in the table. Table 3 lists 150 species of fungi from 47 different aquatic plant species. There seems to be considerable host specificity (with the exception of *Rhizoctonia solani*) reported from 16 different plant species. Quite common were genera *Cercospora* (nine species), *Fusarium* (seven species), and *Pythium* (seven species); all of them are known as cosmopolitan parasites. A common aquatic smut *Doassania*, known to be widespread on aquatic plants, was also reported from several plant species.

114. From the biological control point of view, most research has been done on waterhyacinth. Several fungus species were found to exhibit at least weak pathogenicity to waterhyacinth (see Table 3). Many of them were tested in laboratory or field tests (Charudattan 1973, 1981; Conway and Freeman 1977, Jamil et al. 1983; Lin and Charudattan 1975; Ponnappa 1973; Rahim and Tawfig 1984; Rintz 1973). Often their capacity for infecting a wide range of plants was screened as well. The most promising species were tested in small- and large-scale field tests mostly in the States of Florida and Louisiana.

115. *Cercospora rodmani* has shown good potential as a biological control agent (Freeman et al. 1982). It is an endemic fungus that invades waterhyacinth readily by the way of stomata on leaves and petioles. It causes leaf-spotting with resultant death of severely infected leaves. An epiphytotic can be induced by spraying waterhyacinth with either freshly prepared or formulated inoculum. Once established, the fungus spreads by airborne conidia to plants in adjacent areas. A formulation of *C. rodmani* has been produced commercially by Abbott Laboratories. Agents for potential use in enhancing the germination of spores in the formulation were tested by Pennington and Theriot (1982). *Cercospora piaropi*, which is closely related to *C. rodmani* in both morphology and disease symptomology, was reported as a cause of severe, large-scale damage on waterhyacinth in Lake Conroe, TX (Martyn 1985).

116. In recent years, several research projects have concentrated on biological control of *Hydrilla* and *Myriophyllum*. An isolate of *Fusarium roseum* "Culmorum" isolated from *Stratiotes aloides* near Wageningen, Holland,

was found to be pathogenic to *H. verticillata* (Charudattan et al. 1980, 1984). As the results were very promising, it was decided to test the fungus on *Hydrilla* in a large-scale study. Although the treated plants were more damaged than the controls, a practical level of biocontrol was not achieved. Pennington (1984, 1985) screened numerous isolates of bacterial and fungal epiphytes from *Hydrilla* for production of enzymes that can lyse specific plant components. Several cellulase-producing fungi were capable of damaging *Hydrilla* significantly in test tube assays. The results of aquaria experiments did not confirm the preliminary tests, apparently because the inoculum used for aquarium tests did not have all the necessary additional nutrients or metabolites that were present in the inoculum used for test tube tests. The results also indicated that *Hydrilla* plants might display a different degree of resistance to cellulolytic fungi depending on their previous life history. A similar approach was used by Gunner (1983, 1984), who inoculated *M. spicatum* and *M. heterophyllum* with enzyme-enhanced pectinolytic or cellulolytic microorganisms originally isolated from these species, which significantly accelerated plant necrosis. Among several isolates Gunner tested, a very promising one was the fungus *Mycocleptodiscus terrestris*. Andrews and Hecht (1981) and Andrews et al. (1982) studied the relationship between *M. spicatum* and the fungus *Acremonium curvulum*. *Acremonium* is apparently pathogenic to milfoil when milfoil is grown under conditions of stress.

117. A survey for pathogens of aquatic weeds was done in northern California during 1980 and 1981. Several fungal isolates proved to be pathogenic to their respective hosts (Bernhardt and Duniway 1985, 1986). The most promising for the potential use in biological control was *Pythium carolinianum* pathogenic to *Myriophyllum aquaticum* (syn. *brasiliense*) (Bernhardt and Duniway 1984).

118. Among numerous papers on fungi associated with seaweeds, at least one should be mentioned here. Miller and Jones (1983) described the dependence of traustochytrids on the amount of carbohydrates released by senescing *Fucus* thalli. Traustochytrid numbers were much lower on young and mature host plants than on the senescing ones.

119. Much research has been done on the role of fungi as decomposers of dead organic matter. Over 100 species of higher filamentous fungi have been reported from saltmarsh grasses from the genus *Spartina* (Gessner and Kohlmayer 1976). Twenty isolates of these fungi were surveyed for their ability to

produce degradative enzymes (Gessner 1980). Enzymes capable of degrading cellulose, cellobiose, lipids, pectin, starch, tannic acid, and xylan were detected. One of the fungi occurring most frequently on *Spartina*, *Buergenerula spartinae*, was tested for its polysaccharidase activities. Nine polysaccharidase-type enzymes were detected with xylanase and cellulase being the most active ones (Torzilli and Andrykovitch 1980). No pectic enzyme activity was detected.

120. Besides enzymes, fungi are known to produce antibiotics and toxins. Stevens et al. (1979) analyzed a filtrate of the culture of *A. eichhorniae* growing in potato-dextrose broth. The filtrate was able to induce disease symptoms in waterhyacinth. The main compound of the filtrate was isolated and described as bostrycin, which is an antibiotic against gram-positive bacteria. Bostrycin itself did not show any effect on waterhyacinth, which means that some other constituent in the filtrate, even if in a very low concentration, was biologically active against the plant. Charudattan and Rao (1982) tried to isolate the toxic substance from *A. eichhorniae*. They also isolated 4-deoxybostrycin and isopropylidene derivatives. All four purified products exhibited phytotoxicity against waterhyacinth and may have potential as broad-spectrum herbicides.

121. Rejmankova et al. (1986) found a filtrate from culture of *Pythium myriotylum* to be very effective in killing duckweeds (*L. gibba*). There is a strong possibility that a phytotoxin may be involved in pathogenicity. Phytotoxins have been isolated from several species of *Pythium* (Martin 1964), including *P. myriotylum* (Csinos and Hendrix 1978). Besides being able to produce toxins, several *Pythium* species are also known to be cellulolytic (Deacon 1979). Different *Pythium* species have often been reported as pathogenic not only to freshwater vascular plants but also to filamentous algae (Beck and Erb 1984) and seaweeds (Kazama and Fuller 1970, Aleem 1980).

Viruses

122. Although the term "virus" has been used from about 1890, knowledge of plant viruses has increased markedly in the last few decades mainly as a result of intensified research by both molecular biologists and plant pathologists. Viruses can be defined, according to Stevens (1983), as submicroscopic particles made of one or more pieces of a single species of nucleic acid, RNA

or DNA, surrounded by proteins. These particles replicate alone or in the presence of similar structures, but only in living cells, using at least some of the host cell enzymes. The dimensions of viruses are usually determined by electron microscopy and measured in nanometers (10^{-9} m). Viruses enter and multiply in a wide range of hosts. They have been recorded from numerous plant families, but those from plants of economic importance have been studied most and are best documented. Each virus is named according to the major host with which it is associated and also the symptoms produced in that host. However, a plant virus may invade a number of different plant species, producing different symptoms in each. Nomenclature and classification of plant viruses are described by Smith (1977).

123. Viruses produce visible or otherwise detectable abnormalities in plants. The most obvious symptoms are external and take the form of foliage color changes: mosaicking, mottling, yellowing, ringspot, and local lesions (Stevens 1983). There are many alterations in metabolism in virus-infected plants, e.g., changes in the pattern of carbohydrate accumulation that may be caused by a limited availability of orthophosphate in virus-infected tissues brought about by an accelerated demand for viral nucleic acid synthesis. Orthophosphate deficiency causes a reduction in the movement of triose phosphate from the chloroplast to the cytoplasm and an inhibition of transport of photoassimilates (Misaghi 1982).

124. The spread or transmission of viruses from one plant to another can be mechanical, or an organism called a "vector" can be involved. The following organisms are often recognized as vectors for viruses; insects (namely aphids), mites, nematodes, and fungi.

125. Only a few papers describe viruses in aquatic plants. MacClement and Richards (1956) in their contribution on viruses in wild plants repeatedly observed virus-type infection in *L. minor*, *P. pectinatus*, *P. crispus*, *Ceratophyllum*, and *Nymphaea*. The peak of virus infection was closely connected with the maximum growth of individual species. The frequency of disease found in species sampled in a random survey was about 7 percent for the aquatic species. Pettet and Pettet (1970) reported a massive dieback of *Pistia stratiotes* in the Ibadan area of Western State, Nigeria, associated with a virus infection. The infection seemed to be spread by the aphid *Rhopalosiphum nymphaeae*. The authors apparently did not try to isolate the virus. There were attempts to explain *M. spicatum* decline in the Chesapeake Bay as a result of

virus infection (Bayley et al. 1968). In a later paper, Bayley (1970) stated that she had not been able to isolate any virus from *Myriophyllum* from that area. Stunted plants of alligatorweed (*A. phyloxerooides*) were found in Florida. Flexuous-rod particles 1.717 μ m in diameter were seen in negatively stained leaf extracts. They were similar to the beet yellow group of viruses. Attempts to transfer a pathogen to healthy plants by manual inoculation or by *Apis gossypii* were unsuccessful (Hill and Zettler 1973). Jones (1980) investigated the cause of leaf mottling of *Spartina* and was able to purify and describe a new virus that is distantly serologically related to agropyron mosaic virus. The name "spartina mottle virus" was proposed for this newly recognized virus. There is a short note in the Proceedings of the Conference on Strategies for Aquatic Weed Management (Chestnut 1982) about viruses, or viruslike particles, found in *Hydrilla* and *Ludwigia*, but no more information is given.

126. Several papers describe algal viruses. A virus pathogenic to certain blue-green algae belonging to the genera *Plectonema*, *Phormidium*, and *Lyngbya* has been isolated from a variety of freshwater habitats in Scotland and has been named D-1 (Daft et al. 1970). This virus has similar characteristics to the LPP-1 virus isolated and described in the United States by Safferman and Morris (1964). The virus causes lysis of the host cell and rapid loss of phycobilin and chlorophyll pigments. Review and summary of the state of art in algal virology until 1972 were presented by Brown (1972), who described the virus *Chara corallina* (CCV), which was isolated, characterized, and used to experimentally infect virus-free *Chara* cells (Skotnicki et al. 1976). Van Etten et al. (1985) and Schuster et al. (1986) described several plaque-forming viruses of the green unicellular *Chlorella*-like algae. They were grouped into 11 classes of PBCV-1 viruses. A list of eucaryotic algae in which viruslike particles have been observed by electron microscopy was published by Dodds (1979). Several investigations carried out under field conditions tested the possible utilization of viruses as control of unwanted algae (Jackson and Sladeczek 1970). Algal viruses might serve as "ideal" algicides, but much research still remains to be done.

127. More research must be done before the use of viruses as a potential control tactic for higher aquatic plants can be considered. There is certainly an enormous gap between information on viruses of crop plants where a

lot of genetic engineering has been employed recently (Shewmaker et al. 1985) and that on viruses infecting aquatic plants, which is almost nonexistent.

Nematodes

128. Nematodes are bilaterally symmetric, cylindroid, unsegmented, bisexual, triploblastic organisms comprising a separate phylum *Nematoda*. Nematodes occurring practically everywhere that life can be supported, inhabit a wide variety of environments, and they are the most numerous of all metazoa. They are found in all soils from the polar region to the tropics, whether arid or humid; they are abundant in the bottom sediments of lakes, rivers, and the oceans. The biosphere is carpeted with a layer of nematodes several centimetres thick (Nicholas 1984). All nematodes, however, even those in soil, are more or less aquatic since they require in the active stage a moisture film in which to move and through which to breathe.

129. Worldwide recognition of nematodes as important causal agents of plant diseases did not occur until the middle of this century. The most obvious adaptation to parasitism in phytoparasitic nematodes is their universal possession of a buccal stylet (Zuckerman et al. 1971). The stylet looks like a miniature hypodermic needle. A nematode feeds on the plant by puncturing the cell wall with this stylet. Through the stylet, enzymes are released into the plant cells. Then, as the high-molecular nutrient substances (proteins, carbohydrates, etc.) are degraded to simpler compounds by the action of these enzymes, the nematodes begin to suck up the food through the stylet. An external, extraintestinal digestion takes place. Species of some genera (e.g., *Meloidogyne* and *Heterodera*) were found to produce feeding tubes (Rumpfenhorst 1984). These seem to be common and lasting structures produced at the feeding sites of sedentary root nematodes. Sedentary root nematodes induce the formation of syncytia and giant cell complexes. The feeding tubes, formed by hardening saliva, are connected to the stylet orifice. The feeding tube is assumed to be a duct into which the plant cell secretes special products and through which these secretions are directed into the feeding ampulla. The nematode feeds on the content of this ampulla. The nematode takes in these secreted products rather than common components of cytoplasm.

130. On the basis of their feeding habits, parasitic nematodes are divided into two main groups:

- a. Endoparasites. These enter the plant and feed from inside. Some nematodes move within the tissues and force their way between cells, injuring the tissues as they feed.
- b. Ectoparasites. These do not enter the plant. They feed from outside and rarely penetrate the host tissues.

Plants frequently suffer from more than one disease. Nematodes are notorious partners in plant disease complexes. They act as follows:

- a. Wounding agents. Wounds and punctures provide entry points for other pathogens; once bacteria or fungi are inside, they survive in the tunnels made by the nematodes.
- b. Host modifiers. Digestive enzymes excreted by nematodes change the nutrient contents of the host plant so that it is more palatable not only for nematodes but also for other pathogens as well.
- c. Vectors. The nematode stylet is very small but still big enough for carrying viruses; also, spores can stick to nematode bodies and be transported.

131. The influence of nematodes on fusarium wilts is a well documented and widely occurring example of synergistic interactions between nematodes and fungi enhancing pathogenicity. Mousa and Hague (1985), for example, described an experiment in which fusarium wilt resistant and susceptible cultivars of soybean were exposed to both *Fusarium oxysporum* and *Meloidogyne incognita* (nematode). Both resistant and susceptible cultivars were severely wilted when nematodes were introduced at the same time or about 2 weeks earlier than *Fusarium*. The interaction of the nematode *Criconebella xenoplex* and *Fusarium* on the growth of peaches was studied by Nyczepir and Pusey (1985). When seedlings were treated with *Fusarium* and nematodes, severe symptoms of necrosis were found in the treatment in which nematode inoculation preceded *Fusarium* about 4 weeks. Similar results were obtained by Pelz et al. (1983), who found that the wilt pathogen of the tomato, *F. oxysporum*, is not pathogenic to cucumber in the absence of nematodes (*M. incognita*), but if cucumber plants are simultaneously attacked by nematodes and *Fusarium*, wilt symptoms develop. Waller and Bridge (1984) showed several other examples of the same phenomenon (see Table 26.1 of their paper).

132. Waller and Bridge (1984) stressed that nematode damage does not simply allow easier entry of the fungus into the host; active predisposition of the whole vascular system is also involved (e.g., due to increased production of auxins).

133. Until recently there have not been much data published on nematodes living as commensals or parasites on aquatic plants. Hopper and Meyers (1967) studied the ecology of nematodes in a seagrass community. The microbial microcosm in *S. alterniflora* internodes was described by Gessner et al. (1972); among other microorganisms, the nematodes of the genus *Oncholaimus* were a frequent part of the microcosm. Potential regulating factors of nematodes and other meiofauna in *Spartina* salt marsh in Louisiana were studied by Fleege et al. (1982). Seasonal changes in epiphytic nematodes in a brackish fjord in Finland were described by Jensen (1984).

134. One of the first reports on a freshwater parasitic nematode was on *Aphelenchoides fragariae* collected on several aquatic plant species in Florida (Smart and Esser 1968). Later, Esser et al. (1985) published a survey of phytoparasitic and free-living nematodes associated with aquatic macrophytes in Florida. They detected 25 genera and 38 species of phytoparasitic nematodes and 52 genera of free-living nematodes. The objective of the survey was to find one or more phytoparasitic nematodes that might serve as biological control agents for noxious macrophytes. Among several other phytoparasitic nematodes, *Hirschmanniella caudacrena* was found in very high numbers in the stems and leaves of *C. demersum*. Nematode densities were as high as 58 nematodes/g of fresh plant tissue (Gerber and Smart 1984). Major symptoms of nematode infestation are varying degrees of plant chlorosis, irregular growth in the form of twisted stems, and development of fewer shoots. Laboratory tests have verified that *H. caudacrena* is pathogenic to *Ceratophyllum* (Gerber et al. 1986a). Gerber et al. (1986a) also studied the infestation of *Hydrilla* with parasitic *A. fragariae*. High densities of the nematode were found in *Hydrilla* buds. The nematodes fed ectoparasitically on the meristem, the leaf primordia, the bud scales, and at the base of buds. The apical meristem of many buds was damaged or completely destroyed. *Aphelenchoides* species are active swimmers and are able to move in any direction in the water. This mobility promotes the spread of nematodes to new host plants and could be important concerning the possible use of this nematode for biocontrol of *H. verticillata* (Gerber et al. 1986a).

135. Nematodes occurring in the rhizosphere of aquatic macrophytes, namely *Potamogeton*, are described by Prejs (1977, 1986). She found that tissue damage appearing as differently colored necrotic patches on rhizomes of *Potamogeton* resulted from nematode (mainly *Hirschmanniella*) feeding.

136. Gerber et al. (1986b) prepared a comprehensive catalog of plant parasitic nematodes associated with aquatic and wetland plants. Table 4 shows several selected examples of aquatic plant-nematode associations. The occurrence of nematodes on several aquatic plant species was described also by Gaevskaya (1969). Aquatic plant nematodes do not seem to be very host specific. The most widespread among aquatic plants are the genera *Hirschmanniella* and *Aphelenchoides*. Table 4 includes 35 plant species and 32 nematode genera. *Hirschmanniella* occurred on 17 and *Aphelenchoides* on 11 plant species. *Melioidogyne*, *Dolichodorus*, and *Helicotylenchus* were also quite common; each of them occurred on 8 plant species.

PART IV: SENESCENCE AND PATHOGEN INTERACTION:
ITS ADVANTAGES AND LIMITATIONS

137. Many data exist supporting the idea that senescent plants are more susceptible to pathogen infection than nonsenescent plants. The exact mechanism responsible for this phenomenon is not known. There is probably more than one reason, but, generally, a senescent plant is weaker and less resistant than nonsenescent ones.

138. Senescence proceeds as a natural part of plant development controlled primarily by internal genetic and correlative factors, but senescence can be induced also by unfavorable environmental factors. Table 5 summarizes the possible means of senescence induction and indicates which of them might be used to manipulate the senescence of undesirable aquatic weeds in the field.

139. The environmental factors do not always induce senescence in the whole population of a plant species. Only a few less vigorous plants may become senescent and die while the rest continue growing. The application of a pathogen at the proper moment, i.e., when the population is partially weakened, may greatly enhance its spread. The scheme in Figure 5 shows the possible interactions of induced senescence with pathogens. Nothing, so far, is known about the possibility of inducing pathogenicity in opportunistic microorganisms that would then attack plants and cause their senescence and, eventually, death.




Figure 5. Scheme of interactions between environmentally induced senescence and pathogens

140. More research is needed before the combined effect of senescence and pathogens can be used in practice. It will be necessary to find what is the most practical factor for inducing senescence in a particular species, what type of pathogen is the most effective, at which stage of senescence a pathogen should be applied, etc. Limitations include varying responses of a plant species to the factor inducing senescence, unpredictable interactions of a pathogen with another microorganism present in water, and development of resistance of plants to pathogens.

PART V: CONCLUSIONS AND RECOMMENDATIONS

Conclusions

Senescence

141. The main facts regarding senescence are as follows:
- a. Senescence is an important and not very well-understood stage in the plant life cycle. It is controlled by internal and external (environmental) factors. The precise mechanism of senescence induction is not known, but, in general, the initiation of senescence involves an imbalance in the relative levels of growth hormones; this change in hormone status can be caused by internal factors or environmental stimuli.
 - b. Among the environmental stimuli involved in senescence initiation, the following were reported: light, day length, temperature, mineral nutrients, and pathogens. When pathogens attack plants, it is not always known whether hormonal imbalances are the cause or the result of infection.
 - c. Senescence is accompanied by a change in enzyme activities. Peroxidases, glutamate dehydrogenase, endopeptidases, and some hydrolases have been reported to increase their activities during senescence. Dehydrogenases, ribulosebiphosphate carboxylase, glutamine synthetase, and glutamate synthase generally decrease their activities during senescence.
 - d. An extensive literature search identifies cytokinins as the most generally effective senescence-retarding growth regulators. Abscisic acid (ABA) and ethylene, on the other hand, are known as promoters of senescence. In many cases, cytokinins and ABA interact in a competitive manner.
 - e. Decreases in chlorophyll content and protein amount are most often measured as senescence characteristics. Senescence may start long before changes in these parameters become apparent. No universal marker of senescence has been found to date.

Microorganisms on aquatic plants

142. The review on microorganisms on aquatic plants is concluded as follows:
- a. Aquatic macrophytes often serve as a substratum for epiphytic microorganisms, namely algae, bacteria, and fungi. The amount and diversity of epiphytic microorganisms generally increase with the age of their host mainly due to the increase in organic matter excreted by the aging host plant.
 - b. Microorganisms live mostly as commensals (saprophytes) on their hosts but may become parasitic as the hosts undergo stress conditions or are approaching senescence.

- c. In submersed aquatic plants that usually have a very thin and reduced cuticle, bacteria do not need wounds for entry into plants. Bacteria often invade and degrade epidermal cell walls. Some are known to produce lytic enzymes. Some bacteria were reported as being able to produce antibiotics.
 - d. Fungi are usually regarded as the main decomposers of dead organic matter in aquatic ecosystems. Many species are known to produce enzymes capable of degrading cellulose, pectin, and starch. Besides enzymes, some fungi also produce toxins and antibiotics. The phytotoxicity of fungi has a potential use as bioherbicides. In this study, 150 fungal species were found to be reported as potential parasites on aquatic plants.
 - e. Epiphytic algae do not usually cause any harm to their host plants unless the colonization is so dense that it shades plants or, especially in combination with inorganic silt particles, it becomes so heavy that the host plants are damaged and dragged to the bottom.
 - f. Few viruses were reported from aquatic plants. A potential for biological control may be here, but at this point data are insufficient to even consider this possibility.
 - g. Nematodes are known to contribute to macrophyte degradation in certain cases. They wound the tissue, thus making easy infection entry. So far, at least 32 genera of nematodes parasitic on aquatic plants have been reported.
143. Two main points can be made from this literature search:
- a. Senescence is a process that can be induced by changing the environmental conditions.
 - b. Plants in the senescent stage are more susceptible to pathogens than young or mature but nonsenescent plants.

Recommendations

144. The following research is recommended:
- a. Find the most efficient means of inducing senescence in aquatic weeds that can be eventually applied on a large scale. This would include laboratory tests following the effect of algal epiphytic microflora, light, temperature, nitrogen deficiency, growth regulators (ABA), and sublethal herbicide applications on senescence in selected aquatic plants. These tests should be preceded by another set of experiments aimed at the selection of the most convenient senescence characteristics. Characteristics like chlorophyll and anthocyanin content, protein content, mineral composition of plant biomass, endogenous ABA concentrations, and activities of selected enzymes should be compared in senescing and nonsenescing plants. Within a given aquatic macrophyte species, senescence "markers" should be identified.

- b. Evaluate the effectiveness of a combination of induced senescence with pathogen application. This requires tests on application of pathogens (fungal or bacterial) to aquatic weeds in different stages of senescence, both natural and induced. Attention should be paid to vegetative propagules (turion, tubers) formation in plants with induced senescence. Their susceptibility to pathogens may differ from that of senescent vegetative shoots.
- c. Study the effect of a combination of induced senescence with naturally occurring microorganisms (opportunistic pathogens).
- d. If small-scale research proves that the manipulation of senescence in aquatic weeds following pathogen application is feasible, field tests on a large scale should follow.

REFERENCES

- Adams, M. S., and Prentki, T. R. 1982. "Biology, Metabolism and Functions of Littoral Submersed Weedbeds of Lake Wingra, Wisconsin, USA: A Summary and Review," Archiv fur Hydrobiologie, Supplement, Vol 62, pp 333-409.
- Agami, M., and Waise, I. 1985. "Inter-Relationship Between *Najas marina* L. and Three Other Species of Aquatic Macrophytes," Hydrobiologia, Vol 126, pp 169-173.
- Aiken, S. G., and Walz, K. F. 1979. "Turions of *Myriophyllum exalbescens*," Aquatic Botany, Vol 6, pp 357-363.
- Aleem, A. A. 1980. "*Pythium marinum* Sparrow (Phycomycetes) Infesting *Porphyra leucosticta* Thuret in Mediterranean Sea," Botanica Marina, Vol 23, pp 405-407.
- Allanson, B. R. 1973. "The Fine Structure of the Periphyton on *Chara* sp. and *Potamogeton natans* from Wytham Pond, Oxford, and Its Significance to the Macrophyte-Periphyton Metabolic Model of R. G. Wetzel and H. L. Allen," Freshwater Biology, Vol 3, pp 535-542.
- Allen, H. L. 1971. "Primary Productivity, Chemo-Organotrophy, and Nutritional Interactions of Epiphytic Algae and Bacteria on Macrophytes in the Littoral a Lake," Ecological Monographs, Vol 41, pp 97-127.
- Anderson, L. W. J. 1981. "Control of Aquatic Weeds with Hexazinone," Journal of Aquatic Plant Management, Vol 19, pp 9-14.
- _____. 1986. "Recent Developments and Future Trends in Aquatic Weed Management," Proceedings, European Weed Research Society/AAB 7th Symposium on Aquatic Weeds.
- Andrews, J. H. 1980. "Plant Pathogens as Agents for Biological and Integrated Control of Aquatic Plants," Wisconsin Technical Completion Report, Project Number OWRT A-076-WIS.
- Andrews, J. H., and Hecht, E. P. 1981. "Evidence for Pathogenicity of *Fusarium sporotrichoides* to Eurasian Water Milfoil, *Myriophyllum spicatum*," Canadian Journal of Botany, Vol 59, pp 1069-1077.
- Andrews, J. H., et al. 1982. "Association Between the Fungus *Acremonium curvulum* and Eurasian Water Milfoil, *Myriophyllum spicatum*," Canadian Journal of Botany, Vol 60, pp 1216-1221.
- Anthoni, U., et al. 1980. "Biologically Active Sulphur Compounds from the Green Alga *Chara globularis*," Phytochemistry, Vol 19, pp 1228-1229.
- Aurand, D. 1982. "Nuisance Aquatic Plants and Aquatic Plant Management Programs in the United States," Vol 2, Southeastern Region, MTR-82W47-02, The MITRE Corporation, Metrek Division, VA.
- _____. 1983. "Nuisance Aquatic Plants and Aquatic Plant Management Programs in the United States," Vol 4, Northwestern Region, MTR-82W47-04, The MITRE Corporation, Metrek Division, VA.
- Baker, J. H., and Orr, D. R. 1986. "Distribution of Epiphytic Bacteria on Freshwater Plants," Journal of Ecology, Vol 74, pp 155-165.

- Ballantine, D. L., and Almodovar, L. R. 1977. "The Distribution of Algal Epiphytes on Macrophyte Host, Offshore from La Parguera, Puerto Rico," Proceedings, Association Marine Laboratory Caribbean 12.
- Barko, J. W., and Smart, M. R. 1981. "Comparative Influences of Light and Temperature on the Growth and Metabolism of Selected Submersed Freshwater Macrophytes," Ecological Monographs, Vol 51, pp 219-235.
- Barko, J. W., et al. 1984. "Interactive Influences of Light and Temperature on the Growth and Morphology of Submersed Freshwater Macrophytes," Technical Report A-84-3, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- _____. 1986. "Environmental Factors and Their Consideration in the Management of Submersed Aquatic Vegetation," Journal of Aquatic Plant Management, Vol 24, pp 1-10.
- Batt, T., and Woolhouse, H. W. 1975. "Changing Activities During Senescence and Sites of Synthesis of Photosynthetic Enzymes in Leaves of the Labiate, *Perilla frutescens* (L.) Britt.," Journal of Experimental Botany, Vol 26, pp 569-579.
- Bayley, S. 1970. "The Ecology and Disease of Eurasian Watermilfoil (*Myriophyllum spicatum* L.) in the Chesapeake Bay," Ph.D. thesis, Johns Hopkins University, Baltimore, MD.
- Bayley, S., et al. 1968. "Recent Decline in the Distribution and Abundance of Eurasian Milfoil in Chesapeake Bay," Chesapeake Science, Vol 9, pp 173-181.
- _____. 1978. "Changes in Submerged Aquatic Macrophyte Populations at the Head of Chesapeake Bay, 1958-1975," Estuaries, Vol 1, pp 73-84.
- Bean, G. A., et al. 1973. "Studies on the 'Lake Venice Disease' of Eurasian Milfoil in the Chesapeake Bay," Chesapeake Science, Vol 14, pp 279-280.
- Beck, S. J., and Erb, K. 1984. "Facultative Parasitism of *Spirogyra* sp. (Chlorophyta) by *Saprolegnia asterophora* and *Pythium gracile* (Eumycota Oomycetes)," Journal of Phycology, Vol 20, pp 13-19.
- Bennett, F. D. 1977. "Insects as Agents for Biological Control of Aquatic Weeds," Aquatic Botany, Vol 3, pp 165-174.
- Bernhardt, E. A. 1983. "A Survey for Pathogens of Aquatic Weeds in California for Use in Biological Control," Ph.D. thesis, Department of Plant Pathology, University of California, Davis, CA.
- Bernhardt, E. A., and Duniway, J. M. 1984. "Root and Stem Rot of Parrotfeather (*Myriophyllum brasiliense*)," Plant Disease, Vol 68, pp 999-1003.
- _____. 1985. "A New *Papulospora* Species from the Pondweeds *Potamogeton nodosus* and *P. Crispus* in California," Canadian Journal of Botany, Vol 63, pp 429-431.
- _____. 1986. "Decay of Pondweed and *Hydrilla Hibernacula* by Fungi," Journal of Aquatic Plant Management, Vol 24, pp 20-24.
- Best, E. P. H., et al. 1982. "Studies on Decomposition of *Phragmites australis* Leaves Under Laboratory Conditions," Hydrobiological Bulletin, Vol 16, pp 21-33.

- Birecka, H., et al. 1979. "Perioxidase and Senescence," Journal of Experimental Botany, Vol 30, pp 565-573.
- Biswal, U. C., and Biswal, B. 1984. "Photocontrol of Leaf Senescence," Photochemistry and Photobiology, Vol 39, pp 875-879.
- Blakeman, J. P. 1971. "The Chemical Environment of the Leaf Surface in Relation to Growth of Pathogenic Fungi," Ecology of Leaf Surface Microorganisms, T. F. Preece and C. H. Dickinson, eds., Academic Press, London, pp 255-268.
- Blotnick, J. R., et al. 1980. "Ecological Characteristics of the Rhizosphere Microflora of *Myriophyllum heterophyllum*," Journal of Environmental Quality, Vol 9, pp 207-210.
- Bourn, W. S., and Jenkins, B. 1928. "Rhizoctonia Disease on Certain Aquatic Plants," Botanical Gazette, Vol 85, pp 413-429.
- Bowker, D. W., and Denny, P. 1980. "The Seasonal Succession and Distribution of Epiphytic Algae in the Phyllosphere of *Lemna minor* L.," Archiv fur Hydrobiologie, Vol 90, pp 39-55.
- Boyd, C. E., and Hess, L. W. 1970. "Factors Influencing Shoot Production and Mineral Nutrition Levels in *Typha latifolia*," Ecology, Vol 51, pp 291-301.
- Boylen, C. W., and Brock, T. D. 1974. "A Seasonal Diatom in a Frozen Wisconsin Lake," Journal of Phycology, Vol 10, pp 210-213.
- Brock, T. C. M., et al. 1985. "Field Studies on the Breakdown of *Nuphar lutea* (L.) SM. (*Nymphaeaceae*), and a Comparison of Three Mathematical Models for Organic Weight Loss," Aquatic Botany, Vol 21, pp 1-22.
- Brown, M. R. 1972. "Algal Viruses," Advances in Virus Research, Vol 17, pp 243-277.
- Bulthuis, D. A., and Woelkerling, W. J. 1983. "Biomass Accumulation and Shading Effects of Epiphytes on Leaves of the Seagrass, *Heterozostera tasmanica*, in Victoria, Australia," Aquatic Botany, Vol 16, pp 137-148.
- Butler, R. D., and Simon, E. W. 1971. "Ultrastructural Aspects of Senescence in Plants," Advances in Gerontological Research, Vol 3, pp 73-129.
- Carpenter, S. R. 1980a. "Enrichment of Lake Wingra, WI, by Submersed Macrophyte Decay," Ecology, Vol 65, pp 1145-1155.
- _____. 1980b. "The Decline of *Myriophyllum spicatum* in a Eutrophic Wisconsin Lake," Canadian Journal of Botany, Vol 58, pp 527-535.
- Carter, C. C. 1982. "A Technique for Direct Microscopic Observation of Periphyton Assemblages on Aquatic Macrophytes," Journal of Aquatic Plant Management, Vol 20, pp 53-56.
- Cattaneo, A., and Kalff, J. 1978. "Seasonal Changes in the Epiphyte Community of Natural and Artificial Macrophytes in Lake Memphremagog (Que. and Vt.)," Hydrobiologia, Vol 60, pp 135-144.
- _____. 1980. "The Relative Contribution of Aquatic Macrophytes and Their Epiphytes to the Production of Macrophyte Beds," Limnology and Oceanography, Vol 25, pp 280-289.
- Charudattan, R. 1973. "Pathogenicity of Fungi and Bacteria from India to Hydrilla and Water Hyacinth," Hyacinth Control Journal, Vol 11, pp 44-48.

- Charudattan, R. 1981. "Production, Storage, Germination, and Infectivity of Uredospores of *Ureda eichhorniae* and *Uromyces pontederia*," Phytopathology, Vol 71, pp 1203-1207.
- _____. 1986. "Integrated Control of Waterhyacinth (*Eichhornia crassipes*) with a Pathogen, Insects, and Herbicides," Weed Science, Vol 34, pp 26-30.
- Charudattan, R., and Rao, K. V. 1982. "Bostrycin and 4-Deoxybostrycin: Two Nonspecific Phytotoxins Produced by *Alternaria eichhorniae*," Applied and Environmental Microbiology, Vol 43, pp 846-849.
- Charudattan, R., et al. 1978. "Effects of Fungi and Bacteria on the Decline of Arthropod-Damaged Waterhyacinth (*Eichhornia crassipes*) in Florida," Weed Science, Vol 26, pp 101-107.
- _____. 1980. "Evaluation of *Fusarium roseum* 'Culmorum' as a Biological Control for *Hydrilla verticillata*: Safety," Proceedings, V International Symposium on Biological Control of Weeds, Brisbane, Australia, pp 307-323.
- _____. 1984. "Evaluation of *Fusarium roseum* 'Culmorum' as a Biological Control Agent for *Hydrilla verticillata*," Technical Report A-84-5, prepared by University of Florida, Gainesville, FL, for the US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Chestnut, L. T., ed. 1982. Proceedings of the Conference on Strategies for Aquatic Weed Management, University of Florida, Gainesville, FL.
- Clarkson, D. T., and Hanson, J. B. 1980. "The Mineral Nutrition of Higher Plants," Annual Review of Plant Physiology, Vol 31, pp 239-298.
- Clough, B. F., and Attiwill, P. M. 1980. "Primary Productivity of *Zostera muelleri* Irmisch ex Aschers in Westport Bay (Victoria, Australia)," Aquatic Botany, Vol 9, pp 1-13.
- Cofrancesco, A. F., Jr. 1984. "Ten-Year Update of the Status of Alligatorweed in the Southeast," Miscellaneous Paper A-84-4, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Coler, R. A., and Gunner, H. B. 1969. "The Rhizosphere of an Aquatic Plant (*Lemna minor*)," Canadian Journal of Microbiology, Vol 15, pp 694-696.
- Colquhoun, A. J., and Hillman, J. R. 1972. "The Effect of Abscisic Acid on Senescence in Leaf Discs of Radish, *Raphanus sativus* L.," Planta, Vol 105, pp 213-224.
- Congdon, R. A., and McComb, A. J. 1979. "Productivity of *Ruppia*: Seasonal Changes and Dependence on Light in an Australian Estuary," Aquatic Botany, Vol 6, pp 121-132.
- Connors, I. L. 1967. "An Annotated Index of Plant Diseases in Canada and Fungi Recorded on Plants in Alaska, Canada and Greenland," Publication 1251, Research Branch, Canada Department of Agriculture.
- Conway, K. E. 1978. "A New Species of *Cercospora* from *Limnobium spongia*," Transactions of British Mycological Society, Vol 71, pp 521-523.
- Conway, K. E., and Freeman, T. E. 1977. "Host Specificity of *Cercospora rodmanii*, a Potential Biological Control of Waterhyacinth," Plant Disease Reporter, Vol 61, pp 262-266.

- Conway, K. E., and Kimbrough, J. W. 1975. "A New *Doratomyces* from Water-hyacinth," Mycotaxon, Vol 2, pp 127-131.
- Crowder, A. A., Bristow, J. M., and King, M. R. 1977. "Distribution, Seasonality, and Biomass of Aquatic Macrophytes in Lake Opinicon (Eastern Ontario)," Naturaliste Canadense, Vol 104, pp 441-456.
- Csinos, A., and Hendrix, J. W. 1978. "Parasitic and Nonparasitic Pathogenesis of Tomato Plants by *Pythium myriotylum*," Canadian Journal of Botany, Vol 56, pp 2334-2339.
- Daft, M. J., et al. 1970. "A Virus of Blue-Green Algae from Freshwater Habitats in Scotland," New Phytologist, Vol 69, pp 1029-1038.
- Daly, J. M., and Knoche, H. W. 1976. "Hormonal Involvement in Metabolism of Host-Parasite Interactions," Biochemical Aspects of Plant-Parasite Relationship, J. Friend and D. R. Threlfall, eds., Academic Press, New York, pp 114-133.
- Deacon, J. W. 1979. "Cellulose Decomposition by *Pythium* and Its Relevance to Substrate-Groups of Fungi," Transactions of British Mycological Society, Vol 72, pp 469-477.
- Debusk, T. A., and Ryther, J. H. 1981. "Effect of Seasonality and Plant Density on the Productivity of Some Freshwater Macrophytes," Aquatic Botany, Vol 10, pp 133-142.
- Delbecque, E. J. P. 1983. "A Comparison of the Periphyton of *Nuphar lutea* and *Nymphaea alba*," Developments in Hydrobiology, Vol 17, pp 41-47.
- _____. 1985. "Periphyton on Nymphaeids: An Evaluation of Methods and Separation Techniques," Hydrobiologia, Vol 124, pp 85-93.
- Dodds, J. A. 1979. "Viruses of Marine Algae," Experientia, Vol 35, pp 440-442.
- Drew, E. A. 1979. "Physiological Aspects of Primary Production in Seagrasses," Aquatic Botany, Vol 7, pp 139-150.
- Dykyjova, D. 1978. "Nutrient Uptake by Littoral Communities of Helophytes," Pond Littoral Ecosystems, D. Dykyjova and J. Kvet, eds., Springer-Verlag, Berlin, pp 257-277.
- Esser, R. P., et al. 1985. "A Survey of Phytoparasitic and Free Living Nematodes Associated with Aquatic Macrophytes in Florida," Soil and Crop Science Society of Florida, Vol 44, pp 150-155.
- Fitzgerald, G. P. 1969. "Some Factors in the Competition or Antagonism Among Bacteria, Algae, and Aquatic Weeds," Journal of Phycology, Vol 5, pp 351-359.
- Fleeger, J. W., et al. 1982. "Field Manipulations of Tidal Flushing, Light Exposure and Natant Macrofauna in a Louisiana Salt Marsh: Effect on Meiofauna," Journal of Experimental Marine Biology and Ecology, Vol 56, pp 87-100.
- Fontaine, T. D., and Nigh, D. G. 1983. "Characteristics of Epiphyte Communities on Natural and Artificial Submersed Lotic Plants: Substrate Effects," Archiv fur Hydrobiologie, Vol 96, pp 293-301.

Fowler, M. C. 1977. "Laboratory Trials of a New Triazine Herbicide (DPX 3674) on Various Aquatic Species of Macrophytes and Algae," Weed Research, Vol 17, pp 191-195.

Freeman, T. E. 1977. "Biological Control of Aquatic Weeds with Plant Pathogens," Aquatic Botany, Vol 3, pp 175-184.

_____. 1979. "Control with Plant Pathogens," Proceedings, Research Planning Conference on the Aquatic Plant Control Program A-79-7, US Army Engineer Waterways Experiment Station, Vicksburg, MS, pp 62-67.

Freeman, T. E., et al. 1974. "Utilization of Phytopathogens as Biocontrols for Aquatic Weeds," US Army Engineer Waterways Experiment Station, Vicksburg, MS.

_____. 1981. "Biological Control of Aquatic Plants with Pathogenic Fungi," Technical Report A-81-1, US Army Engineer Waterways Experiment Station, Vicksburg, MS.

_____. 1982. "Rhizoctonia Blight on Waterhyacinth in the United States," Plant Disease, Vol 66, pp 861-862.

Friedrych, J. W., and Huffaker, R. C. 1980. "Photosynthesis, Leaf Resistance, and Ribulose-1,5-Bisphosphate Carboxylase Degradation in Senescing Barley Leaves," Plant Physiology, Vol 65, pp 1103-1107.

Fry, J. C., and Ramsay, A. J. 1977. "Changes in the Activity of Epiphytic Bacteria of *Elodea canadensis* and *Chara vulgaris* Following Treatment with the Herbicide, Paraquat," Limnology and Oceanography, Vol 22, pp 556-562.

Gaevskaya, N. S. 1969. "The Role of Higher Aquatic Plants in the Nutrition of the Animals of Fresh-Water Basins," Boston National Lending Library for Science and Technology, Boston.

Gepstein, S., and Thimann, K. V. 1981. "The Role of Ethylene in the Senescence of Oat Leaves," Plant Physiology, Vol 68, pp 349-354.

Gerber, K., and Smart, G. C. 1984. "Plant Parasitic Nematodes Associated with Aquatic Plants," Proceedings of the First International Congress on Nematology, Guelph, Ontario, Canada, August 5-10, 1984.

Gerber, K., et al. 1986a. "Plant-Parasitic Nematodes Associated with the Aquatic Plants *Ceratophyllum demersum* and *Hydrilla verticillata*," Nematology Circular No. 130, Florida Department of Agriculture and Consumer Service, Gainesville, FL.

_____. 1986b. "A Comprehensive Catalogue of Plant-Parasitic Nematodes Associated with Aquatic and Wetland Plants," Florida Agricultural and Experiment Station, Gainesville, FL.

Gessner, R. V. 1980. "Degradative Enzyme Production by Salt-Marsh Fungi," Botanica Marina, Vol 23, pp 133-139.

Gessner, R. V., and Kohlmayer, J. 1976. "Geographical Distribution and Taxonomy of Fungi from Salt Marsh *Spartina*," Canadian Journal of Botany, Vol 54, pp 2023-2037.

Gessner, R. V., et al. 1972. "The Fungal Microcosm of the Internodes of *Spartina alterniflora*," Marine Biology, Vol 16, pp 269-273.

- Godmaire, H., and Planas, D. 1983. "Potential Effect of *Myriophyllum spicatum* on the Primary Production of Phytoplankton," Developments in Hydrobiology, Vol 17, pp 227-233.
- Godmaire, H., and Planas, D. 1986. "Influence of *Myriophyllum spicatum* on the Species Composition, Biomass and Primary Productivity of Phytoplankton," Aquatic Botany, Vol 23, pp 299-308.
- Godshalk, G., and Wetzel, R. G. 1978. "Decomposition of Aquatic Angiosperms, I. Dissolved Components, II. Particulate Components," Aquatic Botany, Vol 5, pp 281-300, 301-327.
- Goldsborough, L. G., and Robinson, G. G. C. 1985. "Seasonal Succession of Diatom Epiphyton on Dense Mats of *Lemna minor*," Canadian Journal Botany, Vol 63, pp 2332-2339.
- Gough, S. B., and Woelkerling, W. J. 1976. "On the Removal and Quantification of Algal Aufwuchs from Macrophyte Hosts," Hydrobiologia, Vol 48, pp 203-207.
- Grace, J. B., and Tilly, L. J. 1976. "Distribution and Abundance of Submerged Macrophytes, Including *Myriophyllum spicatum* L. (Angiospermae), in a Reactor Cooling Reservoir," Archiv fur Hydrobiologie, Vol 77, pp 475-487.
- Gunner, B. H. 1983. "Microbial Control of Eurasian Watermilfoil," Miscellaneous Paper A-83-4, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- _____. 1984. "Microbial Control of Eurasian Watermilfoil," Proceedings, 18th Annual Meeting, Aquatic Plant Control Research Program, Miscellaneous Paper A-84-4, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Haag, R. W. 1979. "The Ecological Significance of Dormancy in Some Rooted Aquatic Plants," Journal of Ecology, Vol 67, pp 727-738.
- Haag, R. W., and Gorham, P. R. 1977. "Effects of Thermal Effluent on Standing Crop and Net Production of *Elodea canadensis* and Other Submerged Macrophytes in Lake Wabamun, Alberta," Journal of Applied Ecology, Vol 14, pp 835-851.
- Habeshaw, D. 1984. "Effects of Pathogens on Photosynthesis," Plant Diseases, R. K. S. Wood and G. J. Jellis, eds., Blackwell Scientific Publications, Oxford, England, pp 63-72.
- Haller, W. T. 1983. "Hydrilla in Florida," Aquatics, Vol 5, pp 17-18.
- Hannan, H. H., and Dorris, T. C. 1970. "Succession of a Macrophyte Community in a Constant Temperature River," Limnology and Oceanography, Vol 15, pp 442-453.
- Hanson, H. C. 1957. "Dictionary of Ecology," The Catholic University of America, Washington, DC.
- Harlin, M. 1973. "Transfer of Products Between Epiphytic Marine Alga and Host Plant," Journal of Phycology, Vol 9, pp 243-248.
- _____. 1975. "Epiphyte-Host Relations in Seagrass Communities," Aquatic Botany, Vol 1, pp 125-131.

- Harrison, P. G., and Mann, K. H. 1975. "Chemical Changes During the Seasonal Cycle of Growth and Decay in Eelgrass (*Zostera marina* L.) on the Atlantic Coast of Canada," Canadian Fisheries Research Board Journal, Vol 32, pp 615-621.
- Hasler, A. D., and Jones, E. 1949. "Demonstration of the Antagonistic Action of Large Aquatic Plants on Algae and Rotifers," Ecology, Vol 30, pp 359-364.
- Hayslip, H. F., and Zettler, F. W. 1973. "Past and Current Research on Diseases of Watermilfoil (*Myriophyllum spicatum* L.)," Hyacinth Control Journal, Vol 11, pp 38-40.
- Heijs, M. L. F. 1985a. "The Seasonal Distribution and Community Structure of the Epiphytic Algae on *Thalassia hemprichii* (Ehrenb.) Aschers. from Papua New Guinea," Aquatic Botany, Vol 21, pp 295-324.
- Hickman, M., and Klarer, D. M. 1974. "The Growth of Some Epiphytic Algae in a Lake Receiving Thermal Effluent," Archiv Fur Hydrobiologie, Vol 74, pp 403-426.
- Hill, H. R., and Zettler, F. W. 1973. "A Virus-Like Stunting Disease of Alligator Weed from Florida," Phytopathology, Vol 63, p 443, Abstract V.
- Ho, Y. B. 1979. "Chemical Composition Studies on Some Aquatic Macrophytes in Three Scottish Lochs. I. Chlorophyll, Ash, Carbon, Nitrogen and Phosphorus," Hydrobiologia, Vol 63, pp 161-166.
- Hodgson, L. M., and Carter, C. C. 1982. "Effect of Hydrilla Management by Herbicides on a Periphyton Community," Journal of Aquatic Plant Management, Vol 20, pp 17-19.
- Hogestu, K., et al. 1960. "Studies on Antagonistic Relationship Between Phytoplankton and Rooted Aquatic Plants," Japanese Journal of Limnology, Vol 21, pp 124-130.
- Holcomb, G. E. 1982. "Constraints on Disease Development," Biological Control of Weeds with Plant Pathogens, R. Charudattan and H. L. Walker, eds., Wiley, New York, pp 61-71.
- Hopper, B. E., and Meyers, S. P. 1967. "Population Studies on Benthic Nematodes Within a Subtropical Seagrass Community," Marine Biology, Vol 1, pp 85-96.
- Horst, K. R. 1979. Westcott's Plant Disease Handbook, 4th Ed., Van Nostrand Reinhold, New York.
- Hossell, J. C., and Baker, J. H. 1979a. "A Note on the Enumeration of Epiphytic Bacteria by Microscopic Methods with Particular Reference to Two Freshwater Plants," Journal of Applied Bacteriology, Vol 46, pp 87-92.
- _____. 1979b. "Estimation of the Growth Rates of Epiphytic Bacteria and *Lemma minor* in a River," Freshwater Biology, Vol 9, pp 319-327.
- Howard-Williams, C., and Davies, B. R. 1978. "The Influence of Periphyton on the Surface Structure of a *Potamogeton pectinatus* L. Leaf (An Hypothesis)," Aquatic Botany, Vol 5, pp 87-91.
- Howard-Williams, C., and Liport, M. R. M. 1980. "Submerged Macrophyte Communities in a Brackish South African Estuarine-Lake System," Aquatic Botany, Vol 9, pp 101-116.

- Huebert, D. B., and Gorham, P. R. 1983. "Biphasic Mineral Nutrition of the Submersed Macrophyte *Potamogeton pectinatus* L.," Aquatic Botany, Vol 16, pp 269-284.
- Imam, S. H., et al. 1984. "Specificity of Marine Microbial Surface Interactions," Applied and Environmental Microbiology, Vol 48, pp 833-839.
- "Index of Plant Diseases in the United States," US Department of Agriculture Handbook No. 165.
- Jackson, D., and Sladeczek, V. 1970. "Algal Viruses-Eutrophication Control Potential," Yale Scientific Magazine, Vol 44, pp 16-21.
- Jackson, M. B., and Drew, M. C. 1984. "Effect of Flooding on Growth and Metabolism of Herbaceous Plants," Flooding and Plant Growth, T. T. Kozlowski, ed., Academic Press, New York, pp 47-128.
- Jacobs, R. P. W. M., and Noten, T. M. P. A. 1980. "The Annual Pattern of the Diatoms in the Epiphyton of Eelgrass (*Zostera marina* L.) at Roscoff, France," Aquatic Botany, Vol 8, pp 355-370.
- Jamil, K., et al. 1983. "Studies of the Evaluation of Naturally Occurring Fungal Pathogens of Waterhyacinth for Biological Control of the Weed," Indian Journal of Botany, Vol 6, pp 185-189.
- Jana, S., and Choudhuri, M. A. 1979. "Photosynthetic, Photorespiratory, and Respiratory Behaviour of Three Submersed Aquatic Angiosperms," Aquatic Botany, Vol 7, pp 13-19.
- _____. 1980. "Senescence in Submerged Aquatic Angiosperms: Changes in Intact and Isolated Leaves During Aging," New Phytologist, Vol 86, pp 191-198.
- _____. 1981. "Glycolate Metabolism of Three Submersed Aquatic Angiosperms: Effect of Heavy Metals," Aquatic Botany, Vol 11, pp 67-77.
- _____. 1982a. "Glycolate Metabolism of Three Submersed Aquatic Angiosperms During Aging," Aquatic Botany, Vol 12, pp 345-354.
- _____. 1982b. "Characterization of Chilling-Sensitivity of Three Submersed Aquatic Angiosperms," Journal of Aquatic Plant Management, Vol 20, pp 33-36.
- _____. 1982c. "Senescence in Submerged Aquatic Angiosperms: Effects of Heavy Metals," New Phytologist, Vol 90, pp 477-484.
- _____. 1982d. "Changes in the Activities of Ribulose 1,5-Biphosphate and Phosphoenolpyruvate and Carboxylases in Submersed Aquatic Angiosperms During Aging," Plant Physiology, Vol 70, pp 1125-1127.
- _____. 1982e. "Ethylene Production and Senescence in Submerged Aquatic Angiosperms," Aquatic Botany, Vol 13, pp 359-365.
- _____. 1982f. "Glycolate Metabolism of Three Submersed Aquatic Angiosperms During Aging," Aquatic Botany, Vol 12, pp 345-354.
- _____. 1982g. "Changes Occurring During Aging and Senescence in a Submersed Aquatic Angiosperm (*Potamogeton pectinatus*)," Physiologia Plantarum, Vol 55, pp 356-360.
- _____. 1984. "Synergistic Effects of Heavy Metal Pollutions on Senescence in Submersed Aquatic Plants," Water, Air, and Soil Pollution, Vol 21, pp 351-357.

- Jensen, P. 1984. "Ecology of Benthic and Epiphytic Nematodes in Brackish Waters," Hydrobiologia, Vol 108, pp 201-217.
- Johnson, D. A., and King, T. H. 1976. "A Leaf Spot Disease of Three Genera of Aquatic Plants in Minnesota," Plant Disease Reporter, Vol 60, pp 726-730.
- Johnson, D. A., et al. 1976. "A Sclerotium Species Associated with Water Lilies in Minnesota," Plant Disease Reporter, Vol 60, pp 807-808.
- Johnstone, I. M. 1979. "Papua New Guinea Seagrasses and Aspects of the Biology and Growth of *Enhalus acroides* (L.F.) Royle," Aquatic Botany, Vol 7, pp 197-208.
- _____. 1982. "Waterweed Decline: A Resource Allocation Hypothesis," Proceedings, European Weed Research Society, 6th Symposium on Aquatic Weeds, pp 45-53.
- Jones, P. 1980. "Leaf Mottling of *Spartina* Species Caused by a Newly Recognized Virus, *Spartina Mottle Virus*," Annals of Applied Biology, Vol 94, pp 77-81.
- Jones, R. C., and Mayer, K. B. 1983. "Seasonal Changes in the Taxonomic Composition of Epiphytic Algal Communities in Lake Wingra, Wisconsin, U.S.A.," Developments in Hydrobiology, Vol 17, pp 9-16.
- Jones, R. C., et al. 1983. "Phytoplankton as a Factor in the Decline of the Submersed Macrophyte *Myriophyllum spicatum* L. in Lake Wingra, Wisconsin, U.S.A.," Hydrobiologia, Vol 107, pp 213-219.
- Joyner, B. G., and Freeman, T. E. 1973. "Pathogenicity of *Rhizoctonia solani* to Aquatic Plants," Phytopathology, Vol 63, pp 681-685.
- Jupp, B. P., and Spence, D. H. N. 1977. "Limitations on Macrophytes in a Eutrophic Lake, Loch Leven. I. Effect of Phytoplankton," Journal of Ecology, Vol 65, pp 175-186.
- Kandeler, R., et al. 1974. "Genensatzliche Wirkung der Sprossalterung auf die Blütenbildung bei *Lemna paucicostata* und *Lemna gibba*," Biochemie und Physiologie der Pflanzen, Vol 165, pp 331-336.
- Kar, R. K., and Choudhuri, M. A. 1985. "Senescence of *Hydrilla* Leaves in Darkness," Physiologia Plantarum, Vol 63, pp 225-230.
- _____. 1986. "Effect of Light and Spermine on Senescence of *Hydrilla* and Spinach Leaves," Plant Physiology, Vol 80, pp 1030-1033.
- Karpati, V., and Pomogyi, P. 1979. "Accumulation and Release of Nutrients by Aquatic Macrophytes," Symposia Biologica Hungarica, Vol 19, pp 33-42.
- Kazama, F. Y., and Fuller, M. S. 1970. "Ultrastructure of *Porphyra perforata* Infected with *Pythium marinum*, a Marine Fungus," Canadian Journal of Botany, Vol 48, pp 2103-2107.
- Keist, M., and Feller, U. 1984. "Changes in the Pattern of Endopeptidases During Wheat Leaf Senescence," Plant Physiology Supplement 1, Vol 75, p 16.
- Kiorboe, T. 1980b. "Distribution and Production of Submerged Macrophytes in Tipper Grund (Ringkøbing Fjord, Denmark), and the Impact of Waterfowl Grazing," Journal of Applied Ecology, Vol 17, pp 675-687.
- Kirchman, D. L. 1984. "Epiphytic Bacterial Production on *Zostera marina*," Marine Ecology, Vol 15, pp 117-123.

- Kirk, J. J. 1984. "Ability of *Gaeumannomyces graminis* to Benefit from Senescence of the Cereal Root Cortex During Infection," Transactions of British Mycological Society, Vol 82, pp 107-111.
- Kistritz, R. U. 1978. "Recycling of Nutrients in an Enclosed Aquatic Community of Decomposing Macrophytes (*Myriophyllum spicatum*)," Oikos, Vol 30, pp 561-569.
- Klaine, S. J., and Ward, C. H. 1984. "Environmental and Chemical Control of Vegetative Bud Production in *Hydrilla verticillata*," Annals of Botany, Vol 53, pp 503-514.
- Kolattukudy, P. E. 1985. "Enzymatic Penetration of the Plant Cuticle by Fungal Pathogens," Annual Review of Phytopathology, Vol 23, pp 223-250.
- Krishnamani, M. R. S., and Lakshmanan, M. 1976. "Photosynthetic Changes in *Fusarium*-Infected Cotton," Canadian Journal of Botany, Vol 54, pp 1257-1263.
- Kudryavstev, V. M. 1984. "Bacteria on Vascular Aquatic Plants," Gidrobiologicheskij Zhurnal, Vol 19, pp 50-55.
- Kudryavstev, V. M., and Kudryavsteva, N. A. 1980. "Decomposition of *Potamogeton lucens* L. in the Littoral of the Rybinsk Reservoir," Gidrobiologicheskij Zhurnal, Vol 16, pp 14-17.
- Kudryavstev, V. M., and Yeshov, Y. V. 1980. "Abundance Dynamics of Microorganisms in the Decomposition of Macrophytes," Gidrobiologicheskij Zhurnal, Vol 16, pp 20-25.
- Kulshreshtha, M. 1982. "Comparative Ecological Studies on Two Species of *Ceratophyllum*," Proceedings, European Weed Research Society, 6th International Symposium on Aquatic Weeds, Novi Sad, Yugoslavia, pp 29-36.
- Kvet, J. 1975. "Growth and Mineral Nutrients in Shoots of *Typha latifolia* L.," Symposia Biologica Hungarica, Vol 15, pp 113-123.
- Landers, D. H. 1982. "Effects of Naturally Senescing Aquatic Macrophytes on Nutrient Chemistry and Chlorophyll of Surrounding Waters," Limnology and Oceanography, Vol 27, pp 428-439.
- Lauriere, C. 1983. "Enzymes and Leaf Senescence," Physiologia Vegetale, Vol 21, pp 1159-1177.
- Leben, C. 1965. "Epiphytic Organisms in Relation to Plant Disease," Annual Review of Phytopathology, Vol 3, pp 209-230.
- Lee, J. J. 1980. "A Conceptual Model of Marine Detrital Decomposition and the Organisms Associated with the Process," Advances in Aquatic Microbiology, M. R. Droop and H. W. Jannasch, eds., Vol 2, pp 257-291.
- Lemos, M. L., et al. 1985. "Antibiotic Activity of Epiphytic Bacteria Isolated from Intertidal Seaweeds," Microbial Ecology, Vol 11, pp 159-163.
- Leopold, A. C. 1961. "Senescence in Plant Development," Science, Vol 134, pp 1727-1732.
- _____. 1975. "Aging, Senescence, and Turnover in Plants," Bioscience, Vol 25, pp 659-662.
- Leshem, Y. Y. 1986. Processes and Control of Plant Senescence, Elsevier: Amsterdam.

- Lin, C., and Charudattan, R. 1975. "*Sigmoidea* sp., a New Pathogenic Aquatic Hyphomycete of Waterhyacinth," Proceedings of the American Phytopathological Society, Vol 2, p 137.
- Losee, R. F., and Wetzel, R. G. 1983. "Selective Light Attenuation by the Periphyton Complex," Developments in Hydrobiology, Vol 17, pp 89-96.
- Loveless, A. R. 1969. "The Possible Role of Pathogenic Fungi in Local Degeneration of *Salvinia auriculata* Aublet on Lake Kariba," Annals of Applied Biology, Vol 63, pp 61-69.
- Lowe, D. P. 1969. "Check List and Host Index of Bacteria, Fungi, and Mistletoes of British Columbia," Information Report BC-X-32, Forest Research Laboratory, Victoria, British Columbia.
- Lucas, G. B., et al. 1985. Introduction to Plant Diseases, AVI Publication Company, Inc., Westport, CT.
- Lumsden, R. D., et al. 1963. "A Survey of Fungi Associated with Lesioned and Chlorotic Sago Pondweed (*Potamogeton pectinatus*)," Plant Disease Reporter, Vol 47, pp 689-693.
- MacClement, W. D., and Richards, M. G. 1956. "Virus in Wild Plants," Canadian Journal of Botany, Vol 34, pp 793-799.
- Makovetzki, S., and Goldschmidt, E. E. 1976. "A Requirement for Cytoplasmatic Protein Synthesis During Chloroplast Senescence in the Aquatic Plant *Anacharis*," Plant and Cell Physiology, Vol 17, pp 859-862.
- Manos, J. P., and Goldthwaite, J. 1975. "Leaf Tissue Senescence," Plant Physiology, Vol 55, pp 951-953.
- Martin, P. 1964. "Untersuchungen uber ein Phytopathogenes Toxin von *Pythium irregulare* Buisman," Phytopathological Zhurnal, Vol 50, pp 235-249.
- Martyn, R. D. 1985. "Waterhyacinth Decline in Texas Caused by *Cercospora piaropi*," Journal of Aquatic Plant Management, Vol 23, pp 29-31.
- Marvan, P., et al. 1978. "Dynamics of Algal Communities," Pond Littoral Ecosystem, D. Dykyjova and J. Kvet, eds., Springer Verlag, Berlin, pp 314-320.
- Mason, C. F., and Bryant, R. J. 1975. "Production, Nutrient Content, and Decomposition of *Phragmites communis* Trin. and *Typha angustifolia* L.," Journal of Ecology, Vol 63, pp 71-95.
- Menzie, C. A. 1979. "Growth of the Aquatic Plant *Myriophyllum spicatum* L. in a Littoral Area of the Hudson River Estuary," Aquatic Botany, Vol 6, pp 365-375.
- Mickle, A. M., and Wetzel, R. G. 1978. "Effectiveness of Submersed Angiosperm-Epiphyte Complexes on Exchange of Nutrients and Organic Carbon in Littoral Systems. II. Dissolved Organic Carbon," Aquatic Botany, Vol 4, pp 317-329.
- Miller, J. D., and Jones, G. E. B. 1983. "Observations on the Association of Thraustochytrid Marine Fungi with Decaying Seaweed," Botanica Marina, Vol 26, pp 345-351.
- Misaghi, I. J. 1982. Physiology and Biochemistry of Plant-Pathogen Interactions, Plenum Press, New York.

- Moriarty, D. J. W., et al. 1985. "Microbial Biomass and Productivity in Seagrass Beds," Geomicrobiological Journal, Vol 4, pp 21-52.
- Motta, J. J. 1978. "The Occurrence of Fungi on Some Rooted Aquatics from the Chesapeake Bay," Estuaries, Vol 1, pp 101-105.
- Mousa, E. M., and Hague, N. G. M. 1985. "Studies on the Interaction Between *Meloidogyne incognita* and *Fusarium oxysporum glycine* on Soybean," Journal of Nematology, Vol 17, p 506.
- Mulligan, H. F., et al. 1976. "Nitrogen and Phosphorus Fertilization of Aquatic Vascular Plants and Algae in Replicated Ponds. I. Initial Response to Fertilization," Hydrobiologia, Vol 48, pp 101-109.
- Newel, S. Y. 1981. "Fungi and Bacteria in or on Leaves of Eelgrass (*Zostera marina* L.) from Chesapeake Bay," Applied and Environmental Microbiology, Vol 41, pp 1219-1224.
- Nicholas, W. L. 1984. The Biology of Free-Living Nematodes, Clarendon Press, Oxford, England.
- Nichols, D. S., and Keeney, D. R. 1973. "Nitrogen and Phosphorus Release from Decaying Water Milfoil," Hydrobiologia, Vol 42, pp 509-525.
- Nicholson, S. A. 1981. "Changes in Submersed Macrophytes in Chautauque Lake, 1937-1975," Freshwater Biology, Vol 11, pp 523-530.
- Nuttall, P. M. 1985. "Uptake of Phosphorus and Nitrogen by *Myriophyllum aquaticum* (Vellozo) Verd. Growing in a Wastewater Treatment System," Australian Journal of Marine and Freshwater Research, Vol 36, pp 493-507.
- Nuttall, P. M., et al. 1985. "Ammonia Control and Effluent Polishing by Parrot's Feather in an Aquatic Treatment System," Water, Vol 12, pp 17-20.
- Nyczepir, A. P., and Pusey, P. C. 1985. "Interacting of *Criconebella xenoplex* and *Fusarium* sp. on Growth and Feeder Root Necrosis of Peach," Journal of Nematology, Vol 17, p 507.
- Oglesby, R. T., and Vogel, A. 1976. "Changes in Submerged Plants at the South End of Cayuga Lake," Hydrobiologia, Vol 48, pp 251-255.
- Omar, L. S., and Horvath, S. M. 1983. "Elevated Carbon Dioxide Concentrations and Whole Plant Senescence," Ecology, Vol 64, pp 1311-1314.
- O'Neil, Morin J., and Kimball, K. D. 1983. "Relationship of Macrophyte-Mediated Changes in the Water Column to Periphyton Composition and Abundance," Freshwater Biology, Vol 13, pp 403-414.
- Orth, R. J., and Moore, K. A. 1984. "Distribution and Abundance of Submersed Aquatic Vegetation in Chesapeake Bay: An Historical Perspective," Estuaries, Vol 7, pp 531-540.
- Osborne, D. J. 1984. "Ethylene and Plants of Aquatic and Semi-Aquatic Environments: A Review," Plant Growth Regulation, Vol 2, pp 167-185.
- Osborne, D. J., and Cheah, K. S. E. 1982. "Hormones and Foliar Senescence," British Plant Growth Regulator Group Monograph, Vol 8, pp 57-83.
- Panopoulos, N. J., and Peet, R. C. 1985. "The Molecular Genetics of Plant Pathogenic Bacteria and Their Plasmids," Annual Review of Phytopathology, Vol 23, pp 381-419.

- Patience, R. L., et al. 1983. "Changes in Chemical Composition of a Decomposing Aquatic Macrophyte, *Lemna paucicostata*," Journal of Chemical Ecology, Vol 9, pp 889-911.
- Pavgi, M. S., and Singh, L. 1980. "A Physoderma Species Parasitic on Bulrush," Mycopathologia, Vol 71, pp 141-144.
- Pelz, J., et al. 1983. "Einfluss von *Meloidogyne incognita* auf die Wirtseignung der Gurke Gegenüber *Fusarium oxysporum* F. spec. *lycopersici* sowie der Tomate Gegenüber *Fusarium oxysporum* F. spec. *cucumerinum*," Nematologica, Vol 29, pp 443-453.
- Penhale, P. A. 1977. "Macrophyte-Epiphyte Biomass and Productivity in an Eelgrass (*Zostera marina* L.) Community," Journal of Experimental Marine Biology and Ecology, Vol 26, pp 211-224.
- Pennington, J. C. 1984. "Microbiological Control of Hydrilla with Lytic-Enzyme Producing Microorganisms," Proceedings, 18th Annual Meeting, Aquatic Plant Control Research Program, 14-17 November 1983, Raleigh, N.C., Miscellaneous Paper A-84-4, pp 62-65.
- _____. 1985. "Biological Control of *Hydrilla verticillata* (L.f.) Royle with Lytic Enzyme Producing Microorganisms," Technical Report A-85-3, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Pennington, J. C., and Theriot, E. A. 1982. "Compatibility of a *Cercospora rodmanii* Formulation with Selected Herbicide Spray Additives," Miscellaneous Paper A-82-6, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Pettet, A., and Pettet, S. J. 1970. "Biological Control of *Pistia stratiotes* L. in Western State, Nigeria," Nature, Vol 226, p 282.
- Peverly, J. H. 1985. "Element Accumulation and Release by Macrophytes in a Wetland Stream," Journal of Environmental Quality, Vol 14, pp 137-143.
- Phillips, G. L., et al. 1978. "A Mechanism to Account for Macrophyte Decline in Progressively Eutrophicated Freshwaters," Aquatic Botany, Vol 4, pp 103-126.
- Pip, E., and Robinson, G. G. C. 1982. "A Study of the Seasonal Dynamics of Three Phycoperiphytic Communities, Using Nuclear Track Autoradiography. II. Organic Carbon Uptake," Archiv fur Hydrobiologie, Vol 96, pp 47-64.
- Plakidas, A. G. 1936. "Nematodes on Alligator Weed," Plant Disease Reporter, Vol 20, p 22.
- Planas, D., et al. 1981. "Ecological Significance of Phenolic Compounds of *Myriophyllum spicatum*," Verhandlungen der Internationale Vereinigungen der Limnologie, Vol 21, pp 1492-1496.
- Polunin, N. V. C. 1984. "The Decomposition of Emergent Macrophytes in Freshwater," Advances in Ecological Research, Vol 14, pp 115-166.
- Ponnappa, K. M. 1973. "On the Pathogenicity of *Myrothecium roridum* - *Eichhornia crassipes* Isolate," Hyacinth Control Journal, Vol 11, pp 18-20.
- Prejs, K. 1977. "The Nematodes of the Root Region of Aquatic Macrophytes, with Special Consideration of Nematodes Groupings Penetrating the Tissues of Roots and Rhizomes," Ekologia Polska, Vol 25, pp 5-20.

- Prejs, K. 1986. "Nematodes as a Possible Cause of Rhizome Damage in Three Species of *Potamogeton*," Hydrobiologia, Vol 131, pp 281-286.
- Prentki, R. T., et al. 1978. "Nutrient Movements in Lakeshore Marshes," Freshwater Wetlands, R. E. Good et al., eds., Academic Press, New York, pp 169-194.
- Quimby, P. C., and Kay, S. H. 1977. "Hypoxic Quiescence in Alligatorweed," Physiologia Plantarum, Vol 40, pp 163-168.
- Rahim, A. M. A., and Tawfig, S. 1984. "Pathogenicity of Fungi and Bacteria from Sudan to Water Hyacinth," Weeds Research, Vol 24, pp 233-238.
- Ram, H. Y. M., and Kapoor, A. 1976. "In Vitro Culture of Aquatic Weeds and Its Potential Use in Biological Studies," In: "Aquatic Weeds in South East Asia," C. K. Varshney and J. Rzoska, eds., Proceedings of a Regional Seminar on Noxious Aquatic Vegetation, New Delhi, 12-17 December, 1973.
- Ramsay, A. J., and Fry, J. C. 1976. "Response of Epiphytic Bacteria to the Treatment of Two Aquatic Macrophytes with the Herbicide Paraquat," Water Research, Vol 10, pp 453-459.
- Rasmussen, E. 1977. "The Wasting Disease of Eelgrass (*Zostera marina*) and its Effect on Environmental Factors and Fauna," Seagrass Ecosystem. A Scientific Perspective., C. P. McRoy and C. Helfferich, eds., Marcel Dekker, New York, pp 1-51.
- Rejmankova, E. 1973. "Biology of Duckweeds in a Pannonian Fishpond," Symposia Biologica Hungarica, Vol 15, pp 125-131.
- _____. 1979. "The Role of Duckweeds in a Fishpond Ecosystem," Ph.D. dissertation, Czechoslovak Academy of Sciences, Czechoslovakia.
- Rejmankova, E., et al. 1986. "Dynamics of Fungal Infection in Duckweeds," Veroffentlichungen der Beobotanischen Institut, ETH, Stiftung Rubel, Zurich, Vol 87, pp 178-189.
- Renn, C. E. 1936. "Wasting Disease of *Zostera marina*," Biological Bulletin, Vol 70, pp 148-158.
- Riber, H. H., et al. 1983. "Exchange of Phosphorus Between Water, Macrophytes, and Epiphytic Periphyton in the Littoral of Mikolajsie Lake, Poland," Developments in Hydrobiology, Vol 17, pp 235-243.
- Rintz, R. E. 1973. "A Zonal Leaf Spot of Waterhyacinth Caused by *Cephalosporium zonatum*," Hyacinth Control Journal, Vol 11, pp 14.
- Robb, F., et al. 1979. "Cellulolytic Bacteria as Primary Colonizers of *Potamogeton pectinatus* L. (Sago Pond Weed) from a Brackish South-Temperate Coastal Lake," Microbial Ecology, Vol 5, pp 167-177.
- Rogers, K. H., and Breen, C. M. 1980. "Growth and Reproduction of *Potamogeton crispus* in a South African Lake," Journal of Ecology, Vol 68, pp 561-571.
- _____. 1981. "Effects of Epiphyton on *Potamogeton crispus* L. Leaves," Microbial Ecology, Vol 7, pp 351-363.
- _____. 1982. "Decomposition of *Potamogeton crispus* L.: The Effects of Drying on the Pattern of Mass and Nutrient Loss," Aquatic Botany, Vol 12, pp 1-12.

- Rogers, K. H., and Breen, C. M. 1983. "An Investigation of Macrophyte, Epiphyte and Grazer Interactions," Developments in Hydrobiology, Vol 17, pp 217-226.
- Room, P. M. 1986. "Biological Control Is Solving the World's *Salvinia molesta* Problems," Proceedings, European Weed Research Society/AAB 7th Symposium on Aquatic Weeds, pp 271-276.
- Rorslett, B. 1985. "Death of Submerged Macrophytes—Actual Field Observations and Some Implications," Aquatic Botany, Vol 22, pp 7-19.
- Rumpfenhorst, H. J. 1984. "Intracellular Feeding Tubes Associated with Sedentary Plant Parasitic Nematodes," Nematologica, Vol 30, pp 77-85.
- Safferman, R. S., and Morris, M. E. 1964. "Growth Characteristics of the Bluegreen Algal Virus LPP-1," Journal of Bacteriology, Vol 88, pp 771-775.
- Sanders, D. R., and Theriot, E. A. 1985. "Large-Scale Operations Management Test (LSOMT) of Insects and Pathogens for Control for Waterhyacinth in Louisiana; Volume I: Results for 1979-1981," Technical Report A-85-1, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Sand-Jensen, K. 1977. "Effect of Epiphytes on Eelgrass Photosynthesis," Aquatic Botany, Vol 3, pp 55-63.
- _____. 1983. "Physical and Chemical Parameters Regulating Growth of Periphytic Communities," Developments in Hydrobiology, Vol 17, pp 63-71.
- Sasi, P. S., et al. 1979. "Incidence of *Rhizoctonia solani* on *Azolla pinnata*," Current Science, Vol 48, p 216.
- Sastroutomo, S. S. 1982. "The Role of Turions in the Re-Establishment Process of Submerged Weed Vegetation in Japan," Proceedings, European Weed Research Society, 6th Symposium on Aquatic Weeds, pp 77-85.
- Schiemer, F., and Prosser, M. 1976. "Distribution and Biomass of Submerged Macrophytes in Neusiedlersee," Aquatic Botany, Vol 2, pp 289-307.
- Schuster, A. M., et al. 1986. "Characterization of Viruses Infecting a Eukaryotic *Chlorella*-like Green Alga," Virology, Vol 150, pp 170-177.
- Sculthorpe, C. D. 1967. The Biology of Aquatic Vascular Plants, Edward Arnold, London.
- Sharma, K. P. 1985. "Allelopathic Influence of Algae on the Growth of *Eichhornia crassipes* (Mart.) Solms," Aquatic Botany, Vol 22, pp 71-78.
- Shewmaker, C. K., et al. 1985. "Transcription of Cauliflower Mosaic Virus Integrated into Plant Genomes," Virology, Vol 140, pp 281-288.
- Siddiqui, I. A., et al. 1973. "Distribution of Plant Parasitic Nematodes in California," State of California Department of Food and Agriculture.
- Skotnicki, A., et al. 1976. "Further Study on *Chara corallina* Virus," Virology, Vol 75, pp 457-468.
- Sladeckova, A. 1962. "Limnological Investigation Methods for the Periphyton ("Aufwuchs") Community," Botanical Review, Vol 28, pp 286-350.
- Smart, G. C., Jr., and Esser, R. P. 1968. "*Aphelenchoides fragariae* in Aquatic Plants," Plant Disease Reporter, Vol 52, p 455.

- Smedegaard-Petersen, V. 1985. "The Limiting Effect of Disease Resistance on Yield," Annual Review of Phytopathology, Vol 23, pp 475-490.
- Smith, H., and Grierson, D., eds. 1982. "The Molecular Biology of Plant Development," Botanical Monographs, Vol 18, pp 257-281.
- Smith, K. M. 1977. Plant Viruses, Wiley, New York.
- Solski, A. 1962. "Mineralization of Aquatic Plants," Polskie Archiwum Hydrobiologii, Vol 10, pp 167-196. (In Polish with English summary).
- Sondergaard, M. 1983. "Heterotrophic Utilization and Decomposition of Extracellular Carbon Released by the Aquatic Angiosperm *Littorella uniflora* (L.) Aschers.," Aquatic Botany, Vol 16, pp 59-73.
- Sparrow, F. K. 1974. "Observations on Chytridiaceous Parasites of Phanerogams. XIX. A Physoderma on Eurasian Water Milfoil (*Myriophyllum spicatum*)," American Journal of Botany, Vol 61, pp 174-180.
- Spence, D. H. N. 1972. "Light on Freshwater Macrophytes," Transactions of Botanical Society, Edinburgh, Vol 41, pp 491-505.
- Spencer, D. F., and Anderson, L. W. J. 1986a. "Photoperiod Response in Monoecious and Dioecious *Hydrilla verticillata*," Weed Science, Vol 34, pp 551-557.
- _____. 1986b. "Influence of Photoperiod on Growth, Pigment Composition, and Vegetative Propagule Formation for *Potamogeton nodosus* Poir. and *Potamogeton pectinatus* L."
- Sterry, P. R., et al. 1985. "Changes in the Concentrations of Short-Chain Carboxylic Acids and Gases During Decomposition of the Aquatic Macrophytes, *Lemna paucicostata* and *Ceratophyllum demersum*," Freshwater Biology, Vol 15, pp 139-153.
- Stevens, K. L., et al. 1979. "The Antibiotic Bostrycin from *Alternaria eichhorniae*," Water Chemistry, Vol 18, pp 1579-1580.
- Stevens, W. A. 1983. "Virology of Flowering Plants," Chapman and Hall, New York.
- Stowe, C. W. 1982. "Diatoms Epiphytic on the Emergent Grass *Spartina alterniflora* in a Louisiana Salt Marsh," Transactions of American Microscopical Society, Vol 101, pp 162-173.
- Strobel, G. A. 1977. "Bacterial Phytotoxins," Annual Review of Microbiology, Vol 31, pp 205-224.
- Strother, S. 1984. "Control of Phosphatase Activities of Duckweed by Senescence and Phosphorus Deficiency," Biologia Plantarum, Vol 26, pp 309-311.
- Taylor, D. I. 1983. "The Effects of a Major Macrophyte Regression upon Primary Production in the Littoral of Swartvlei," Archiv fur Hydrobiologie, Vol 96, pp 345-353.
- Thomas, H., and Stoddart, J. L. 1980. "Leaf Senescence," Annual Review of Plant Physiology, Vol 31, pp 83-111.
- Thomas, S. M., et al. 1978. "Ribulose 1,5-Biphosphate Carboxylase/Oxygenase Activity and Photorespiration During the Aging of Flag Leaves of Wheat," Journal of Experimental Botany, Vol 29, pp 1161-1168.

- Titus, J. E., and Stone, W. H. 1982. "Photosynthetic Response of Two Submerged Macrophytes to Dissolved Inorganic Carbon Concentration and pH," Limnology and Oceanography, Vol 27, pp 151-160.
- Torzilli, A. P., and Andrykovitch, G. 1980. "Cell Wall Degrading Enzymes Produced by the Salt Marsh Fungus *Buergenerula spartinae*," Botanica Marina, Vol 23, pp 645-650.
- Trudeau, P. N. 1982. "Nuisance Aquatic Plants and Aquatic Plant Management Programs in the United States," Vol 3, Northeastern and North Central Region, MTR-82W47-03, The MITRE Corporation, Metrek Division, VA.
- Tutin, T. G. 1938. "The Autecology of *Zostera marina* in Relation to its Wasting Disease," New Phytologist, Vol 37, pp 50-71.
- Tyndall, R. W. 1982. "Nuisance Aquatic Plants and Aquatic Plant Management Programs in the United States," Vol 1, Southwestern Region, MTR-82W47-01, The MITRE Corporation, Metrek Division, VA.
- Tyndall, R. W., et al. 1981. "Integrated Aquatic Weed Management," MTR-81W87, The MITRE Corporation, Metrek Division, VA.
- Unni, S. K. 1976. "Production of Submerged Aquatic Plant Communities of Doodhadhari Lake Raipur, (M.P. India)," Hydrobiologia, Vol 48, pp 175-177.
- Van der Velde, G. 1978. "Structure and Functioning of a Nymphaeid-Dominated System," Proceedings, European Weed Research Society, 5th Symposium on Aquatic Weeds, pp 127-133.
- Van Etten, J. L., et al. 1985. "Lytic Viruses Infecting a *Chlorella*-like Alga," Virology, Vol 140, pp 135-143.
- Van Montfrans, J., et al. 1982. "Preliminary Studies of Grazing by *Bittium varium* on Eelgrass Periphyton," Aquatic Botany, Vol 14, pp 75-89.
- Van Vierssen, W., and Prins, T. C. 1985. "On the Relationship Between the Growth of Algae and Aquatic Macrophytes in Brackish Water," Aquatic Botany, Vol 21, pp 165-179.
- Van Zon, J. C. J. 1977. "Introduction to Biological Control of Aquatic Weeds," Aquatic Botany, Vol 3, pp 105-110.
- Varghese, G., and Singh, G. 1973. "Progress in the Search for Natural Enemies of *Hydrilla* in Malaysia," In: "Aquatic Weeds in South East Asia," C. K. Varshney and J. Rzoska, eds., Proceedings of a Regional Seminar on Noxious Aquatic Vegetation, New Delhi, 12-17 December 1973.
- Verhoeven, J. T. A. 1975. "*Ruppia* Communities in the Camargue, France. Distribution and Structure in Relation to Salinity and Salinity Fluctuations," Aquatic Botany, Vol 1, pp 217-241.
- _____. 1980. "The Ecology of *Ruppia*-Dominated Communities in Western Europe III. Aspects of Production, Consumption and Decomposition," Aquatic Botany, Vol 8, pp 209-253.
- Wagner, D. T. 1969. "A Monocentric, Holocarpic Fungus in *Lemna minor* L.," Nova Hedwigia, Vol 18, pp 203-208.
- Waite, T. D., and Kurucz, C. 1977. "The Kinetics of Bacteria Degradation of the Aquatic Weed *Hydrilla* sp.," Water, Air, and Soil Pollution, Vol 7, pp 33-43.

- Waite, T. D., and Mitchell, R. 1976. "Some Benevolent and Antagonistic Relationship Between *Ulva lactuca* and Its Microflora," Aquatic Botany, Vol 2, pp 13-22.
- Waller, J. M., and Bridge, J. 1984. "Effects of Pathogen Interactions on Tropical Crop Production," Plant Disease: Infection, Damage and Loss, R. K. S. Weed and G. L. Jellis, eds., Blackwell Scientific Publications, Oxford, England.
- Wareing, P. F., and Phillips, I. D. J. 1985. Growth and Differentiation in Plants, Pergamon Press, New York.
- Weber, J. A., and Nooden, L. D. 1974. "Turion Formation and Germination in *Myriophyllum verticillatum*: Phenology and Interpretation," The Michigan Botanist, Vol 13, pp 151-165.
- _____. 1976. "Environmental and Hormonal Control of Turion Formation in *Myriophyllum verticillatum*," Plant and Cell Physiology, Vol 17, pp 721-731.
- Weitzel, R. L., ed. 1979. "Methods and Measurements of Periphyton Communities: A Review," American Society for Testing and Materials, Philadelphia, PA.
- Westlake, D. F. 1965. "Some Basic Data for Investigations of the Productivity of Aquatic Macrophytes," In: "Primary Productivity in Aquatic Environments," C. R. Goldman, ed., Proceedings of IBP Symposium Pallanza, Italy, pp 231-248.
- Wetzel, R. G. 1983. "Attached Algal-Substrata Interactions: Fact or Myth, and When and How," Developments in Hydrobiology, Vol 17, pp 207-215.
- Wetzel, R. G., and Allen, H. L. 1970. "Functions and Interactions of Dissolved Organic Matter and the Littoral Zone in Lake Metabolism and Eutrophication," Productivity Problems of Freshwaters, Z. Kajak and A. Hillbricht-Ilkowska, eds., Warsaw, PWN Polish Scientific Publishers, pp 333-347.
- Wetzel, R. G., and Penhale, P. A. 1979. "Transport of Carbon and Excretion of Dissolved Organic Carbon by Leaves and Roots/Rhizomes in Seagrasses and Their Epiphytes," Aquatic Botany, Vol 6, pp 149-158.
- Willcocks, A. P. 1982. "Colonization and Distribution of Red Algal Epiphytes *Melobesia medioeris* and *Smithora naiadum* on the Seagrass *Phyllospadix Torreyi*," Aquatic Botany, Vol 12, pp 365-373.
- Woolhouse, H. W., and Jenkins, G. I. 1983. "Physiological Responses, Metabolic Changes and Regulation During Leaf Senescence," Growth and Functioning of Leaves, J. E. Dale and F. L. Milthorpe, eds., Cambridge University Press, pp 449-487.
- Yeo, R. R., and Thurston, J. R. 1984. "The Effect of Dwarf Spikerush (*Eleocharis coloradoensis*) on Several Submersed Aquatic Weeds," Journal of Aquatic Plant Management, Vol 22, pp 52-56.
- Yoshida, Y. 1961. "Chloroplast Senescence in Detached *Elodea* Leaves," Protoplasma, Vol 54, pp 476-492.
- _____. 1970. "Control of Chloroplast Senescence of Detached *Elodea* Leaves," Botanical Magazine, Tokyo, Vol 83, pp 137-143.

Zettler, F. W., and Freeman, T. E. 1972. "Plant Pathogens as Biocontrols of Aquatic Plants," Annual Review of Phytopathology, Vol 10, pp 455-470.

Zuckerman, B. M., et al., ed. 1971. Plant Parasitic Nematodes, Academic Press, New York.

BIBLIOGRAPHY

- Agami, M., et al. 1980. "Growth and Photosynthesis of *Najas marina* L. as Affected by Light Intensity," Aquatic Botany, Vol 9, pp 285-289.
- Barltrop, J., et al. 1982. "Response of *Hydrilla* to Selective Dyes," Journal of Environmental Science and Health, Vol A17, pp 725-735.
- Best, E. P. H., and Meulemans, J. T. 1978. "Primary Productivity in the Submerged Aquatic Macrophyte *Ceratophyllum demersum*," Verhandlungen Internationale Verein Limnologie, Vol 20, pp 88-93.
- _____. 1979. "Photosynthesis in Relation to Growth and Dormancy in *Ceratophyllum demersum*," Aquatic Botany, Vol 6, pp 53-65.
- Brock, T. C. M., et al. 1983. "Structure and Annual Biomass Production of *Nymphoides peltata* (Gmel.) O. Kuntze (*Menyanthaceae*)," Aquatic Botany, Vol 17, pp 167-186.
- Catling, P. M., and Dobson, I. 1985. "The Biology of Canadian Weeds. 69. *Potamogeton crispus* L.," Canadian Journal of Plant Science, Vol 65, pp 655-668.
- Charudattan, R., and Lin, C. V. 1973. "Isolates of *Penicillium*, *Aspergillus*, and *Trichoderma* Toxic to Aquatic Plants," Hyacinth Control Journal, Vol 11, pp 70-73.
- Charudattan, R., and Walker, H. L., eds. 1982. Biological Control of Weeds with Plant Pathogens, Wiley, New York.
- Chitwood, B. G., and Allen, M. W. 1959. "Nemata," Freshwater Biology, W. T. Edmonson, ed., Wiley, New York, pp 368-401.
- Conway, K. E. 1975. "*Cercospora rodmani*, a New Pathogen of Water Hyacinth with Biological Control Potential," Canadian Journal of Botany, Vol 54, pp 1079-1083.
- Cunningham, J. J., et al. 1984. "Temporal Response of the Macrophyte, *Potamogeton perfoliatus* L., and Its Associated Autotrophic Community to Atrazine Exposure in Estuarine Microcosms," Estuaries, Vol 7, pp 519-530.
- Dawson, F. H. 1983. "Practical Applications of a Shading Material for Macrophyte Control in Water Courses," Aquatic Botany, Vol 17, pp 301-308.
- Donnermayer, G. N., and Smart, M. M. 1985. "The Biomass and Nutritive Potential of *Vallisneria americana* Michx in Navigation Pool 9 of the Upper Mississippi River," Aquatic Botany, Vol 22, pp 33-44.
- Dooris, P. M., et al. 1982. "Effect of Hydrilla-Growth Inhibiting Extracts upon the Growth of *Scenedesmus obliquus* Growth Inhibition," Journal of Environmental Science and Health, Vol A17, pp 639-645.
- Elser, H. J. 1969. "Observation on the Decline of the Water Milfoil and Other Aquatic Plants, Maryland, 1962-1967," Hyacinth Control Journal, Vol 8, pp 52-60.

- Esteves, F. A. 1979. "Die Bedeutung der aquatischen Makrophyten für den Stoffhaushalt des Schöhsees. I. Die Produktion an Biomasse, II. Die Organische Hauptbestandteile und der Energiegehalt der Aquatischen Makrophyten," Archiv für Hydrobiologie/Supplement, Vol 57, pp 117-187.
- Gallagher, J. C. 1978. "Decomposition Processes: Summary and Recommendations," Freshwater Wetlands. Ecological Processes and Management Potential, R. E. Good et al., ed., Academic Press, New York, pp 145-151.
- Gessner, R. V. 1976. "In Vitro Growth and Nutrition of *Buergenerula spartinae*, a Fungus Associated with *Spartina alterniflora*," Mycologia, Vol 68, pp 583-599.
- Gons, H. J. 1982. "Structural and Functional Characteristics of Epiphyton and Epipelon in Relation to Their Distribution in Lake Vechten," Hydrobiologia, Vol 95, pp 79-114.
- Gophen, M. 1982. "Unusually Dense Watermilfoil (*Myriophyllum spicatum* L.) Vegetation in the Southern Basin of Lake Kinneret (Israel) in 1979," Aquatic Botany, Vol 13, pp 307-315.
- Grace, J. B., and Wetzel, R. G. 1978. "The Production of Biology of Eurasian Watermilfoil (*Myriophyllum spicatum* L.): A Review," Journal of Aquatic Plant Management, Vol 16, pp 1-11.
- Heijs, M. L. F. 1985b. "Some Structural and Functional Aspects of the Epiphytic Component of Four Seagrass Species (*Cymadaceoideae*) from Papua New Guinea," Aquatic Botany, Vol 23, pp 225-247.
- Hodgson, L. M., and Linda, S. B. 1984. "Response of Periphyton and Phytoplankton to Chemical Control of *Hydrilla* in Artificial Pools," Journal of Aquatic Plant Management, Vol 22, pp 48-51.
- Hootsmans, M. J. M., and Vermaat, J. E. 1985. "The Effect of Periphyton Grazing by Three Epifaunal Species on the Growth of *Zostera marina* L. under Experimental Conditions," Aquatic Botany, Vol 22, pp 83-88.
- "Index of Plant Virus Diseases." 1966. US Department of Agriculture Handbook No. 307.
- Jones, R. C., and Adams, M. S. 1982. "Seasonal Variations in Photosynthetic Response of Algae Epiphytic on *Myriophyllum spicatum* L.," Aquatic Botany, Vol 13, pp 317-330.
- Kiorboe, T. 1980. "Production of *Ruppia cirrhosa* (Petagna) Grande in Mixed Beds in Ringkøbing Fjord (Denmark)," Aquatic Botany, Vol 9, pp 135-143.
- Lammens, E. H. R. R., and Van der Velde, G. 1978. "Observations on the Decomposition of *Nymphoides peltata* (Gmel.). Kuntze (Menyanthaceae) with Special Regard to the Leaves," Aquatic Botany, Vol 4, pp 331-346.
- Marshall, B. E., and Junor, F. J. R. 1981. "The Decline of *Salvinia molesta* on Lake Kariba," Hydrobiologia, Vol 83, pp 477-484.
- Martyn, R. D., and Freeman, T. E. 1978. "Evaluation of *Acremonium zonatum* as a Potential Biocontrol Agent of Waterhyacinth," Plant Disease Reporter, Vol 62, pp 604-608.
- Mickle, A. M., and Wetzel, R. G. 1978. "Effectiveness of Submersed Angiosperm-Epiphyte Complexes on Exchange of Nutrients and Organic Carbon in Littoral Systems. I. Inorganic Nutrients," Aquatic Botany, Vol 4, pp 303-316.

- Mickle, A. M., and Wetzel, R. G. 1979. "Effectiveness of Submersed Angiosperm-Epiphyte Complexes on Exchange of Nutrients and Organic Carbon in Littoral Systems. III. Refractory Organic Carbon," Aquatic Botany, Vol 6, pp 339-355.
- Myuge, S. G. 1964. "Plant-Parasitic Nematodes," Kolos, Moskva, Israel Program for Scientific Translations, Jerusalem, 1969.
- Newell, S. Y., and Fell, J. W. 1980. "Mycoflora on Turtlegrass (*Thalassia testudium* Konig) as Recorded After Seawater Incubation," Botanica Marina, Vol 23, pp 265-275.
- Newton, R. J. 1977. "Abscisic Acid Effects on Fronds and Roots of *Lemma minor* L.," American Journal of Botany, Vol 64, pp 45-49.
- Norton, D. C. 1978. "Ecology of Plant-Parasitic Nematodes," Wiley, New York.
- Ostrowski, M. L., and Zettler, E. R. 1986. "Chemical Defenses in Aquatic Plants," Journal of Ecology, Vol 74, pp 279-287.
- Park, D. 1972. "On the Ecology of Heterotrophic Micro-organisms in Fresh Water," Transactions of British Mycological Society, Vol 58, pp 291-299.
- Prentki, R. T., et al. 1979. "The Role of Submersed Waterbeds in Internal Loading and Interception of Allochthonous Materials in Lake Wingra, Wisconsin, USA," Archiv fur Hydrobiologie, Vol 57, pp 221-250.
- Rho, J., and Gunner, H. B. 1978. "Microfloral Response to Aquatic Weed Decomposition," Water Research, Vol 12, pp 165-170.
- Robertson, K. P., and Woolhouse, H. W. 1984. "Studies on the Seasonal Course of Carbon Uptake of *Eriophorum vaginatum* in a Moorland Habitat. I. Leaf Production and Senescence, II. The Seasonal Course of Photosynthesis," Journal of Ecology, Vol 72, pp 423-435, 685-700.
- Sand-Jensen, K., and Sondergaard, M. 1981. "Phytoplankton and Epiphyte Development and Their Shading Effect on Submerged Macrophytes in Lakes of Different Nutrient Status," Internationale Revue der Gesamte Hydrobiologie, Vol 66, pp 529-552.
- Schatz, S. 1984. "Degradation of *Lamionaria saccharina* by Saprobic Fungi," Mycologia, Vol 76, pp 426-432.
- Stanley, R. A., et al. 1976. "Effect of Season and Water Depth on Eurasian Watermilfoil," Journal of Aquatic Plant Management, Vol 14, pp 32-36.
- Stowe, C. W., and Gosselink, J. G. 1985. "Metabolic Activity of the Epiphytic Community Associated with *Spartina alterniflora*," Gulf Research Report, Vol 8, pp 21-25.
- Suberkropp, K., and Arsuffi, T. L. 1984. "Degradation, Growth, and Changes in Palatability of Leaves Colonized by Six Aquatic Hyphomycete Species," Mycologia, Vol 76, pp 398-407.
- Suge, H., and Kusanagi, T. 1975. "Ethylene and Carbon Dioxide: Regulation of Growth in Two Perennial Aquatic Plants, Arrowhead and Pondweed," Plant and Cell Physiology, Vol 16, pp 65-72.
- Tillberg, E., et al. 1979. "Growth Cycles in *Lemma gibba* Cultures and Their Effect on Growth Rate and Ultrastructure," Physiologia Plantarum, Vol 46, pp 5-12.

Vanderkloet, S. 1977. "Distribution, Seasonality, and Biomass of Aquatic Macrophytes in Lake Opinicon (Eastern Ontario)," Naturaliste Canadien, Vol 104, pp 441-456.

Wang, T. L., and Woolhouse, H. W. 1982. "Hormonal Aspects of Senescence in Plant Development," British Plant Growth Regulator Group, Monograph No. 8, pp 5-25.

Wasfy, E. H., et al. 1984. "Changes in Peroxidase Isoenzyme Pattern of Bean Hypocotyl Due to Infection with *Rhizoctonia solani*," Transactions of British Mycological Society, Vol 82, pp 154-156.

Whitfield, A. K. 1984. "The Effect of Prolonged Aquatic Macrophyte Senescence on the Biology of the Dominant Fish Species in a Southern African Coastal Lake," Estuarine, Coastal and Shelf Science, Vol 18, pp 315-319.

Table 1

List of Epiphytic Bacteria Associated with Aquatic Plants

Plant Species	Type of Bacteria	Reference
<i>Equisetum</i> sp.	<i>Azotobacter</i>	Kudryavstev (1984)
<i>Hydrilla verticillata</i>	"Gram negative, nonmotile rods"	Waite and Kurucz (1977)
	<i>Pseudomonas</i>	Pennington (1985) ↓
	<i>Acinetobacter</i>	
	<i>Aeromonas</i>	
	<i>Flavobacterium</i>	
	<i>Bacillus</i>	
	<i>Enterobacter</i>	
	<i>Alcaligena</i>	
	<i>Flexibacter</i>	
	<i>Erwinia</i>	
	<i>Myconostoc</i>	
	<i>Rhodopseudomonas</i>	
	<i>Serratia</i>	
	<i>Caulobacter</i>	
	<i>Agrobacterium</i>	
	<i>Hafnia</i>	
	<i>Klebsiella</i>	
	<i>Sporocytophaga</i>	
<i>Lemna paucicostata</i>	"Rod-shaped, coccoid bacteria"	Patience et al. (1983)
<i>Myriophyllum heterophyllum</i>	"Gram negative rods, pseudomonad type"	Gunner (1983)
<i>Myriophyllum spicatum</i>	<i>Heterophyllum</i> sp.	Gunner (1983)
	<i>Pectinolytic isolate</i> Br-2	Gunner (1983)
	<i>Cyanoacterial consortium</i>	Gunner (1983)
<i>Najas flexillis</i>	<i>Caulobacter</i> sp.	Allen (1971)
	<i>Pseudomonas</i> sp.	Allen (1971)
<i>Potamogeton pectinatus</i>	<i>Bacillus</i> sp.	Howard-Williams and Davies (1978)
	<i>Caulobacterium</i> sp.	Robb et al. (1979)
	"Rough and smooth walled bacilli short rods, gram positive"	Robb et al. (1979)
<i>Potamogeton</i> sp.	<i>Azotobacter</i>	Kudryavstev (1984)
<i>Sagittaria</i> sp.	<i>Azotobacter</i>	Kudryavstev (1984)

(Continued)

Table 1 (Concluded)

Plant Species	Type of Bacteria	Reference
<i>Spartina alterniflora</i>	<i>Metallogenium</i>	Gessner et al. (1972)
	<i>Hyphomicrobium</i>	Gessner et al. (1972)
	<i>Mycoplasma</i>	Gessner et al. (1972)
<i>Typha angustifolia</i>	<i>Cytophaga</i> group	Mason and Bryant (1975)
	<i>Pseudomonas fluorescens</i>	Mason and Bryant (1975)
	<i>Pseudomonas marginalis</i>	Mason and Bryant (1975)
	<i>Erwinia</i> sp.	Mason and Bryant (1975)
<i>Zostera marina</i>	"Long rods"	Newel (1981)
"Intertidal seaweeds"	<i>Pseudomonas-Alteromonas</i> group	Lemos et al. (1985)

Table 2

Fungi Reported as Potential Pathogens from Aquatic Plants

Fungus	Host Plant	Reference
Plasmodiophorales (Plasmodiophoromycetes)		
<i>Ligneria junci</i> <i>Tetramyxa parasitica</i>	<i>Alisma plantago-aquatica</i> <i>Ruppia cirrhosa</i>	Index of Plant Diseases Verhoeven (1975)
Labyrinthulales (Labyrinthulomycetes)		
<i>Labyrinthula</i> sp.	<i>Zostera marina</i>	Renn (1936)
Chytridiales (Chytridiomycetes)		
<i>Cladochytrium maculare</i> <i>Reesia amoeboides</i>	<i>Alisma plantago-aquatica</i> <i>Lemna minor</i>	Index of Plant Diseases Wagner (1969)
Blastocladales (Chytridiomycetes)		
<i>Physoderma</i> sp. <i>Physoderma thilumalachari</i>	<i>Myriophyllum spicatum</i> <i>Scirpus articulatus</i>	Sparrow (1974) Pavgi and Singh (1980)
Lagenidiales (Oomycetes)		
<i>Lagenidium muenschari</i>	<i>Potamogeton</i> sp.	Index of Plant Diseases
Saprolegniales (Oomycetes)		
<i>Saprolegnia asterophora</i>	<i>Spirogyra</i> sp.	Beck and Erb (1984)
Peronosporales (Oomycetes)		
<i>Albugo bliti</i>	<i>Alternanthera phylloxeroides</i>	Index of Plant Diseases

(Continued)

(Sheet 1 of 10)

Table 2 (Continued)

Fungus	Host Plant	Reference
Peronosporales		
(Oomycetes)		
(Continued)		
<i>Pythium afertile</i>	<i>Elodea canadensis</i>	Bernhardt (1983)
<i>Pythium afertile</i>	<i>Myriophyllum spicatum</i>	↓
<i>Pythium afertile</i>	<i>Potamogeton crispus</i>	
<i>Pythium afertile</i>	<i>Potamogeton pectinatus</i>	
<i>Pythium carolinianum</i>	<i>Myriophyllum brasiliense</i>	
<i>Pythium carolinianum</i>	<i>Myriophyllum spicatum</i>	
<i>Pythium debaryanum</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Pythium ultimum</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Pythium gracile</i>	<i>Spirogyra</i> sp.	Beck and Erb (1984)
<i>Pythium helicoides</i>	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Pythium myriotylum</i>	<i>Lemna gibba</i>	Rejmankova et al. (1986)
<i>Pythium myriotylum</i>	<i>Lemna minor</i>	Rejmankova et al. (1986)
<i>Pythium myriotylum</i>	<i>Spirodela Polyrhiza</i>	Rejmankova et al. (1986)
<i>Pythium</i> sp.	<i>Eichhornia crassipes</i>	Charudattan (1973)
<i>Pythium</i> sp.	<i>Hydrilla verticillata</i>	Varghese and Singh (1973)
<i>Pythium</i> sp.	<i>Nuphar luteum</i>	Van der Velde (1978)
<i>Pythium</i> sp.	<i>Nymphaea odorata</i>	Index of Plant Diseases
<i>Pythium</i> sp.	<i>Nymphaea tuberosa</i>	Index of Plant Diseases
<i>Pythium</i> sp.	<i>Potamogeton nodosus</i>	Bernhardt (1983)
<i>Pythium</i> sp.	<i>Potamogeton pectinatus</i>	Lumsden et al. (1963)
<i>Pythium</i> sp.	<i>Potamogeton perfoliatus</i>	Motta (1978)
<i>Pythium</i> sp.	<i>Ruppia maritima</i>	Motta (1978)
<i>Pytophthora parasitica</i>	<i>Hydrilla verticillata</i>	Freeman (1977)
<i>Pythiogeton autossytum</i>	<i>Typha latifolia</i>	Index of Plant Diseases
Helotiales		
(Ascomycotina)		
<i>Pezizella oenontherae</i>	<i>Lythrum salicaria</i>	Index of Plant Diseases

(Continued)

(Sheet 2 of 10)

Table 2 (Continued)

Fungus	Host Plant	Reference
Hypocreales		
(Ascomycotina)		
<i>Claviceps nigricans</i>	<i>Eleocharis acicularis</i>	Connors (1967)
<i>Claviceps purpurea</i>	<i>Phragmites communis</i>	Index of Plant Diseases
Rhytismatales		
(Ascomycotina)		
<i>Lophodermium typhinum</i>	<i>Typha latifolia</i>	Index of Plant Diseases
Sphaeriales		
(Ascomycotina)		
<i>Buergenerula spartinae</i>	<i>Spartina alterniflora</i>	Torzilli and Andrykovitch (1980)
<i>Chaetomium funiculum</i>	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Coniochaeta</i> sp.	<i>Heteranthera dubia</i>	Bernhardt (1983)
<i>Guignardia cepalariae</i>	<i>Altermanthera phyloxeroides</i>	Index of Plant Diseases
<i>Hendersonia arundinacea</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Hendersonia typhae</i>	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Leptosphaeria culmorum</i>	<i>Eleocharis pervula</i>	Connors (1967)
<i>Leptosphaeria eustoma</i>	<i>Typha latifolia</i>	Lowe (1969)
<i>Leptosphaeria typharum</i>	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Lulworthia</i> sp.	<i>Zostera marina</i>	Newel (1981)
<i>Mycosphaerella pontederiae</i>	<i>Nuphar advena</i>	Index of Plant Diseases
<i>Mycosphaerella pontederiae</i>	<i>Nymphaea odorata</i>	Index of Plant Diseases
<i>Mycosphaerella typhae</i>	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Mycosphaerella</i> sp.	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Ophiobolus halimus</i>	<i>Zostera marina</i>	Newel (1981)
<i>Ophiobolus</i> sp.	<i>Typha latifolia</i>	Horst (1979)
<i>Pleospora graminis</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Pleospora typhae</i>	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Sphaerulina pedicellata</i>	<i>Spartina alterniflora</i>	Gessner et al. (1972)

(Continued)

(Sheet 3 of 10)

Table 2 (Continued)

Fungus	Host Plant	Reference
Uredinales		
(Urediniomycetes)		
<i>Nyssopsora echinata</i>	<i>Oenante sarmentosa</i>	Conners (1967)
<i>Puccinia eleocharidis</i>	<i>Eleocharis palustris</i>	Conners (1967)
<i>Puccinia mogrusiana</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Puccinia phragmitis</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Puccinia scirpi</i>	<i>Numphoides peltata</i>	Munda and Garrasi (1978)
<i>Uredo eichhorniae</i>	<i>Eichhornia crassipes</i>	Freeman et al. (1981)
<i>Uredo maculans</i>	<i>Alternanthera phylloxeroides</i>	Index of Plant Diseases
<i>Uromyces acuminatus</i>	<i>Spartina alterniflora</i>	Conners (1967)
<i>Uromyces pontederiae</i>	<i>Pontederia lanceolata</i>	Charudattan (1981)
Ustilaginales		
(Ustilaginomycete)		
<i>Burrillia decipiens</i>	<i>Nymphaea cordata</i>	Conners (1967)
<i>Burrillia limnanthemii</i>	<i>Nymphaea cordata</i>	Conners (1967)
<i>Burrillia pustulata</i>	<i>Sagittaria</i> sp.	Conners (1967)
<i>Doassansia alismatis</i>	<i>Alisma plantago-aquatica</i>	Index of Plant Diseases
<i>Doassansia alismatis</i>	<i>Alisma geyeri</i>	Index of Plant Diseases
<i>Doassansia deformans</i>	<i>Sagittaria</i> sp.	Conners (1967)
<i>Doassansia furva</i>	<i>Sagittaria</i> sp.	Conners (1967)
<i>Doassansia hydrophila</i>	<i>Potamogeton natans</i>	Lowe (1969)
<i>Doassansia intermedia</i>	<i>Sagittaria</i> sp.	Conners (1967)
<i>Doassansia martanoffiana</i>	<i>Potamogeton nodosus</i>	Conners (1967)
<i>Doassansia obscura</i>	<i>Sagittaria</i> sp.	Conners (1967)
<i>Doassansia oculata</i>	<i>Potamogeton</i> sp.	Index of Plant Diseases
<i>Doassansia opaca</i>	<i>Sagittaria</i> sp.	Conners (1967)
<i>Doassansia sagittariae</i>	<i>Sagittaria</i> sp.	Conners (1967)

(Continued)

(Sheet 4 of 10)

Table 2 (Continued)

Fungus	Host Plant	Reference	
Ustilaginales			
(Ustilaginomycete)			
(Continued)			
<i>Entyloma nymphaeae</i>	<i>Nuphar advena</i>	Index of Plant Diseases	
<i>Entyloma nymphaeae</i>	<i>Nymphaea odorata</i>	Index of Plant Diseases	
<i>Entyloma nymphaeae</i>	<i>Nymphaea tuberosa</i>	Index of Plant Diseases	
<i>Neovossia iowensis</i>	<i>Phragmites communis</i>	Index of Plant Diseases	
<i>Tracya lemnae</i>	<i>Spirodela polyrrhiza</i>	Connors (1967)	
Melanconiales			
(Coelomycetes)			
<i>Cryptomela typhae</i>	<i>Typha latifolia</i>	Index of Plant Diseases	
<i>Gleosporium nymphaeearum</i>	<i>Nymphaea tuberosa</i>	Connors (1967)	
<i>Marssonina</i> sp.	<i>Sagittaria</i> sp.	Connors (1967)	
Sphaeropsidales			
(Coelomycetes)			
<i>Apicarpella</i> sp.	<i>Eichhornia crassipes</i>	Freeman et al. (1974)	
<i>Chaetophoma</i> sp.	<i>Hydrilla verticillata</i>	Charudattan (1973)	
<i>Coniothyrium</i> sp.	<i>Ceratophyllum</i> sp.	Bernhardt (1983)	
<i>Coniothyrium</i> sp.	<i>Potamogeton crispus</i>	Bernhardt (1983)	
<i>Hymenopsis hydrophila</i>	<i>Typha latifolia</i>	Index of Plant Diseases	
<i>Phoma sorghina</i>	<i>Eichhornia crassipes</i>	Rahim and Tawfig (1984)	
<i>Phoma</i> sp.	<i>Myriophyllum spicatum</i>	Bernhardt (1983)	
<i>Phoma</i> sp.	<i>Potamogeton pectinatus</i>	Lumsden et al. (1963)	
<i>Phyllosticta fatisscens</i>	<i>Nuphar advena</i>	Index of Plant Diseases	
<i>Phyllosticta fatisscens</i>	<i>Nuphar odorata</i>	↓	
<i>Phyllosticta fatisscens</i>	<i>Nuphar advena</i>		
<i>Phyllosticta nymphaeacea</i>			
<i>Phyllosticta typhina</i>	<i>Typha latifolia</i>		
<i>Selenophoma graminis</i>	<i>Phragmites communis</i>		
<i>Septoria lythrina</i>	<i>Lythrum salicaria</i>		
<i>Septoria oenanthis</i>	<i>Oenante sarmentosa</i>		Connors (1967)
<i>Septoria punctoides</i>	<i>Eleocharis pauciflora</i>		Connors (1967)
<i>Septoria villarsiae</i>	<i>Nymphoides peltate</i>		Munda and Garrasi (1978)

(Continued)

(Sheet 5 of 10)

Table 2 (Continued)

Fungus	Host Plant	Reference
Sphaeropsidales (Coelomycetes) (Continued)		
<i>Stagnospora typhiodearum</i>	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Stagnospora</i> sp.	<i>Phragmites communis</i>	Index of Plant Diseases
Agnomycetales = Mycelia sterilia (Hyphomycetes)		
<i>Papulaspora aspera</i>	<i>Potamogeton crispus</i>	Bernhardt and Duniway (1985)
<i>Papulaspora aspera</i>	<i>Potamogeton nodosus</i>	Bernhardt and Duniway (1985)
<i>Papulaspora</i> sp.	<i>Potamogeton perfoliatus</i>	Motta (1978)
<i>Papulaspora</i> sp.	<i>Ruppia maritima</i>	Motta (1978)
<i>Rhizoctonia solani</i>	<i>Altermanthera phyloxerooides</i>	Freeman et al. (1974)
	<i>Azolla pinnata</i>	Sasi et al. (1979)
	<i>Hydrilla verticillata</i>	Joyner and Freeman (1973)
	<i>Hydrocotyle umbellata</i>	Joyner and Freeman (1973)
	<i>Lemna minor</i>	Joyner and Freeman (1973)
	<i>Limnobiium spongia</i>	Joyner and Freeman (1973)
	<i>Myriophyllum aquaticum</i>	Joyner and Freeman (1973)
	<i>Myriophyllum spicatum</i>	Freeman et al. (1981)
	<i>Najas flexilis</i>	Bourn and Jenkins (1928)
	<i>Pistia stratiotes</i>	Joyner and Freeman (1973)
	<i>Pontederia lanceolata</i>	Joyner and Freeman (1973)
	<i>Potamogeton pectinatus</i>	Bourn and Jenkins (1928)
	<i>Potamogeton perfoliatus</i>	Bourn and Jenkins (1928)
	<i>Ruppia maritima</i>	Bourn and Jenkins (1928)
	<i>Salvinia rotundifolia</i>	Joyner and Freeman (1973)
	<i>Vallisneria spiralis</i>	Bourn and Jenkins (1928)

(Continued)

(Sheet 6 of 10)

Table 2 (Continued)

Fungus	Host Plant	Reference
Agnomycetales		
= <i>Mycelia sterilia</i>		
(Hyphomycetes)		
(Continued)		
<i>Rhizoctonia</i> sp.	<i>Eichhornia crassipes</i>	Freeman (1977)
<i>Sclerotium</i> <i>hydrophyllum</i>	<i>Myriophyllum spicatum</i>	Hayslip and Zettler (1973)
<i>Sclerotium rolfsii</i>	<i>Hydrilla verticillata</i>	Freeman 1977
<i>Sclerotium</i> sp.	<i>Nuphar luteum</i>	Johnson et al. (1976)
<i>Sclerotium</i> sp.	<i>Nymphaea odorata</i>	Johnson et al. (1976)
Hyphomycetales		
= Moniliales		
(Hyphomycetes)		
<i>Acremonium curbulum</i>	<i>Myriophyllum spicatum</i>	Andrews et al. (1982)
<i>Acremonium zonatum</i>	<i>Eichhornia crassipes</i>	Freeman (1977)
<i>Acremonium</i> sp.	<i>Zostera marina</i>	Newel (1981)
<i>Alternaria</i> <i>altermantherae</i>	<i>Altermanthera phyloxeroidea</i>	Holcomb (1982)
<i>Alternaria</i> <i>eichhorniae</i>	<i>Eichhornia crassipes</i>	Sharma (1985)
<i>Alternaria</i> <i>tenuissima</i>	<i>Elodea canadensis</i>	Bernhardt (1983)
<i>Alternaria</i> <i>tenuissima</i>	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Alternaria</i> sp.	<i>Nymphaea odorata</i>	Horst (1979)
<i>Alternaria</i> sp.	<i>Salvinia auriculata</i>	Loveless (1969)
<i>Anquillospora</i> <i>crassa</i>	<i>Potamogeton pectinatus</i>	Bernhardt (1983)
<i>Aspergillus awomori</i>	<i>Hydrilla verticillata</i>	Pennington (1985)
<i>Aspergillus</i> sp.	<i>Eichhornia crassipes</i>	Charudattan (1973)
<i>Aspergillus</i> sp.	<i>Hydrilla verticillata</i>	Freeman (1977)
<i>Bipolaris</i> <i>stenospila</i>	<i>Eichhornia crassipes</i>	Freeman (1977)
<i>Cephalosporium</i> sp.	<i>Eichhornia crassipes</i>	Rahim and Tawfig (1984)
<i>Cephalosporium</i> sp.	<i>Hydrilla verticillata</i>	Charudattan (1973)
<i>Cephalosporium</i> sp.	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Cephalosporium</i> sp.	<i>Potamogeton perfoliatus</i>	Motta (1978)
<i>Cephalosporium</i> sp.	<i>Potamogeton pectinatus</i>	Bernhardt (1983)
<i>Cephalosporium</i> sp.	<i>Ruppia maritima</i>	Motta (1978)
<i>Cercospora</i> <i>altermantherae</i>	<i>Altermanthera phyloxeroidea</i>	Index of Plant Diseases
<i>Cercospora</i> <i>alismatis</i>	<i>Alisma plantago-aquatica</i>	Index of Plant Diseases

(Continued)

(Sheet 7 of 10)

Table 2 (Continued)

Fungus	Host Plant	Reference
Hyphomycetales = Moniliales (Hyphomycetes) (Continued)		
<i>Cercospora alismatis</i>	<i>Sagittaria</i> sp.	Horst (1979)
<i>Cercospora exotica</i>	<i>Nymphaea odorata</i>	Horst (1979)
<i>Cercospora limnobiai</i>	<i>Limnobia spongia</i>	Conway (1978)
<i>Cercospora nymphaeacea</i>	<i>Nuphar luteum</i>	Charudattan et al.
<i>Cercospora nymphaeacea</i>	<i>Nymphaea odorata</i>	Index of Plant Diseases
<i>Cercospora piaropi</i>	<i>Eichhornia crassipes</i>	Freeman (1977)
<i>Cercospora pontederiae</i>	<i>Pontederia cordata</i>	Connors (1967)
<i>Cercospora rodmani</i>	<i>Eichhornia crassipes</i>	Conway (1978)
<i>Cercospora sagittariae</i>	<i>Sagittaria</i> sp.	Connors (1967)
<i>Cladosporium</i> sp.	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Cladosporium</i> sp.	<i>Zostera marina</i>	Newel (1981)
<i>Curvularia lunata</i>	<i>Eichhornia crassipes</i>	Rahim and Tawfig (1984)
<i>Curvularia</i> sp.	<i>Hydrilla verticillata</i>	Varghese and Singh (1973)
<i>Curvularia</i> sp.	<i>Potamogeton nodosus</i>	Bernhardt (1983)
<i>Curvularia</i> sp.	<i>Potamogeton pectinatus</i>	Lumsden et al. (1963)
<i>Cylindrocarpon</i> sp.	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Dactylella microaquatica</i>	<i>Myriophyllum spicatum</i>	Hayslip and Zettler (1973)
<i>Dactylella aquatica</i>	<i>Potamogeton crispus</i>	Bernhardt (1983)
<i>Dichotomophthoropsis numphaearum</i>	<i>Brasenia schreberi</i>	Johnson and King (1976)
	<i>Nuphar luteum</i>	Johnson and King (1976)
<i>Didymaria alismatis</i>	<i>Nymphaea orodata</i>	Johnson and King (1976)
	<i>Nymphaea tuberosa</i>	Johnson and King (1976)
<i>Didymaria alismatis</i>	<i>Alisma plantago-aquatica</i>	Index of Plant Diseases
<i>Didymaria alismatis</i>	<i>Alisma geyeri</i>	Index of Plant Diseases
<i>Didymaria alismatis</i>	<i>Sagittaria</i> sp.	Connors (1967)
<i>Diplosporium</i> sp.	<i>Potamogeton pectinatus</i>	Bernhardt (1983)
<i>Doratomyces eichhornius</i>	<i>Eichhornia crassipes</i>	Conway and Kimbrough (1975)
<i>Drechslera spicifera</i>	<i>Eichhornia crassipes</i>	Rahim and Tawfig (1984)
<i>Epicoccum nigrum</i>	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Flagellospora curvula</i>	<i>Myriophyllum spicatum</i>	Bernhardt (1983)

(Continued)

(Sheet 8 of 10)

Table 2 (Continued)

Fungus	Host Plant	Reference
Hyphomycetales = Moniliales (Hyphomycetes) (Continued)		
<i>Flagellospora curvula</i>	<i>Potamogeton pectinatus</i>	Bernhardt (1983)
<i>Flagellospora stricta</i>	<i>Myriophyllum spicatum</i>	Hayslip and Zettler (1973)
<i>Fusarium acuminatum</i>	<i>Sagittaria</i> sp.	Conners (1967)
<i>Fusarium equiseti</i>	<i>Eichhornia crassipes</i>	Rahim and Tawfig (1984)
<i>Fusarium lateritium</i>	<i>Heteranthera dubia</i>	Bernhardt (1983)
<i>Fusarium moniliforme</i>	<i>Eichhornia crassipes</i>	Rahim and Tawfig (1984)
<i>Fusarium moniliforme</i>	<i>Myriophyllum spicatum</i>	Hayslip and Zettler (1973)
<i>Fusarium moniliforme</i>	<i>Potamogeton pectinatus</i>	Bernhardt (1983)
<i>Fusarium nivale</i>	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Fusarium oxysporium</i>	<i>Myriophyllum spicatum</i>	Hayslip and Zettler (1973)
<i>Fusarium roseum</i>	<i>Eichhornia crassipes</i>	Freeman et al. (1982)
<i>Fusarium roseum</i>	<i>Hydrilla verticillata</i>	Charudattan et al. (1984)
<i>Fusarium roseum</i>	<i>Myriophyllum spicatum</i>	Hayslip and Zettler (1973)
<i>Fusarium solani</i>	<i>Eichhornia crassipes</i>	Jamil et al. (1983)
<i>Fusarium solani</i>	<i>Hydrilla verticillata</i>	Charudattan et al. (1978)
<i>Fusarium solani</i>	<i>Myriophyllum spicatum</i>	Hayslip and Zettler (1973)
<i>Fusarium sporotrichoides</i>	<i>Myriophyllum spicatum</i>	Andrews and Hecht (1981)
<i>Fusarium tricinctum</i>	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Geotrichium</i> sp.	<i>Potamogeton crispus</i>	Bernhardt (1983)
<i>Gliocephalis</i> sp.	<i>Hydrilla verticillata</i>	Charudattan (1973)
<i>Gliocladium roseum</i>	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Heterosporium maculatum</i>	<i>Scirpus</i> sp.	Lowe (1969)
<i>Heterosporium maculatum</i>	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Humicola</i> sp.	<i>Hydrilla verticillata</i>	Pennington (1985)
<i>Hyaloflorae</i> sp.	<i>Potamogeton pectinatus</i>	Lumsden et al. (1963)
<i>Mycoleptodiscus terrestris</i>	<i>Eichhornia crassipes</i>	Horst (1979)

(Continued)

(Sheet 9 of 10)

Table 2 (Concluded)

Fungus	Host Plant	Reference
Hyphomycetales = Moniliales (Hyphomycetes) (Continued)		
<i>Mycoleptodiscus terrestris</i>	<i>Myriophyllum spicatum</i>	Gunner (1983)
<i>Myrothecium roridum</i>	<i>Eichhornia crassipes</i>	Freeman et al. (1982)
<i>Napicladium arundinaceum</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Nigrospora oryzae</i>	<i>Potamogeton pectinatus</i>	Bernhardt (1983)
<i>Nigrospora</i> sp.	<i>Eichhornia crassipes</i>	Charudattan (1973)
<i>Oidiodendron</i> sp.	<i>Hydrilla verticillata</i>	Charudattan (1973)
<i>Ovularia numphaerum</i>	<i>Numphaea odorata</i>	Index of Plant Diseases
<i>Paecilomyces</i> sp.	<i>Ceratophyllum</i> sp.	Bernhardt (1983)
<i>Papularia arundinis</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Penicillium</i> sp.	<i>Hydrilla verticillata</i>	Freeman (1977)
<i>Periconia</i> sp.	<i>Hydrilla verticillata</i>	Charudattan (1973)
<i>Pullaria pullulans</i>	<i>Potamogeton pectinatus</i>	Lumsden et al. (1963)
<i>Ramularia aquatilis</i>	<i>Potamogeton</i> sp.	Index of Plant Diseases
<i>Rhynchosporium alismatis</i>	<i>Alisma plantago-aquatica</i>	Connors (1967)
<i>Rhynchosporium alismatis</i>	<i>Sagittaria</i> sp.	Connors (1967)
<i>Scolecotrichum graminis</i>	<i>Phragmites communis</i>	Index of Plant Diseases
<i>Scolecotrichum typhae</i>	<i>Typha latifolia</i>	Index of Plant Diseases
<i>Spicariopsis</i> sp.	<i>Salvinia auriculata</i>	Loveless (1969)
<i>Sitgmina</i> sp.	<i>Hydrilla verticillata</i>	Charudattan (1973)
<i>Sigmoidea</i> sp.	<i>Eichhornia crassipes</i>	Lin and Charudattan (1975)
<i>Sporobolomyces roseus</i>	<i>Nuphar advena</i>	Connors (1967)
<i>Trichoderma viride</i>	<i>Potamogeton perfoliatus</i>	Motta (1978)
<i>Trichoderma viride</i>	<i>Ruppia maritima</i>	Motta (1978)
<i>Trichoderma</i> sp.	<i>Eichhornia crassipes</i>	Charudattan (1973)
<i>Trichoderma</i> sp.	<i>Hydrilla verticillata</i>	Freeman (1977)
<i>Trichoderma</i> sp.	<i>Myriophyllum spicatum</i>	Bernhardt (1983)
<i>Trichoderma</i> sp.	<i>Potamogeton crispus</i>	Bernhardt (1983)
<i>Verticillium</i> sp.	<i>Hydrilla verticillata</i>	Varghese and Singh (1973)

Table 3

Aquatic Plants from Which the Pathogenic Fungi Were Reported

<u>Plant Species</u>	<u>Fungus</u>	<u>Reference</u>
<i>Alisma plantago-aquatica</i>	<i>Cercospora alismatis</i>	Index of Plant Diseases
	<i>Cladochytrium maculare</i>	
	<i>Didymaria alismatis</i>	
	<i>Doassansia alismatis</i>	
	<i>Rhynchosporium alismatis</i>	Conners (1967)
<i>Alisma geyeri</i>	<i>Didymaria alismatis</i>	Index of Plant Diseases
	<i>Doassansia alismatis</i>	Index of Plant Diseases
<i>Alternanthera philoxeroides</i>	<i>Albugo bliti</i>	Index of Plant Diseases
	<i>Alternaria alternantherae</i>	Holcomb and Antonopoulos (1976)
	<i>Cercospora alternantherae</i>	Index of Plant Diseases
	<i>Guignardia cepalariae</i>	Index of Plant Diseases
	<i>Rhizoctonia solani</i>	Freeman et al. (1974)
	<i>Uredo maculans</i>	Index of Plant Diseases
<i>Azolla pinnata</i>	<i>Phizoctonia solani</i>	Sasi et al. (1979)
<i>Brasenia schreberi</i>	<i>Dichotomophthoropsis numphaearum</i>	Johnson and King (1976)
<i>Ceratophyllum demersum</i>	<i>Coniothyrium</i> sp.	Bernhardt (1983)
	<i>Paecilomyces</i> sp.	Bernhardt (1983)
<i>Eichhornia crassipes</i>	<i>Acremonium zonatum</i>	Freeman (1977)
	<i>Alternaria eichhorniae</i>	Sharma (1985)
	<i>Apicarpella</i> sp.	Freeman et al. (1974)
	<i>Aspergillus</i> sp.	Charudattan (1973)
	<i>Bipolaris stenospila</i>	Freeman (1977)
	<i>Cephalosporium</i> sp.	Rahim and Tawfig (1984)
	<i>Cercospora piaropi</i>	Freeman (1977)
	<i>Cercospora rodmanii</i>	Conway (1975)
	<i>Curvularia lunata</i>	Rahim and Tawfig (1984)
	<i>Doratomyces eichhornius</i>	Conway and Kimbrough (1975)
	<i>Drechslera spicifera</i>	Rahim and Tawfig (1984)
	<i>Fusarium equiseti</i>	Rahim and Tawfig (1984)
	<i>Fusarium moniliforme</i>	Rahim and Tawfig (1984)
	<i>Fusarium solani</i>	Jamil et al. (1983)
	<i>Fusarium roseum</i>	Freeman et al. (1978)
<i>Myrothecium roridum</i>	Freeman et al. (1978)	

(Continued)

(Sheet 1 of 7)

Table 3 (Continued)

Plant Species	Fungus	Reference
<i>Eichhornia crassipes</i> (Continued)	<i>Mycoleptodiscus terrestris</i>	Horst (1979)
	<i>Nigrospora</i> sp.	Charudattan (1973)
	<i>Phoma sorphina</i>	Rahim and Tawfig (1984)
	<i>Pythium</i> sp.	Charudattan (1973)
	<i>Rhizoctonia</i> sp.	Freeman (1977)
	<i>Sigmoidea</i> sp.	Lin and Charudattan (1975)
	<i>Uredo eichhorniae</i>	Freeman et al. (1981)
	<i>Trichoderma</i> sp.	Freeman et al. (1981)
<i>Eleocharis acicularis</i>	<i>Claviceps nigricans</i>	Connors (1967)
<i>Eleocharis palustris</i>	<i>Puccinia eleocharidis</i>	Connors (1967)
<i>Eleocharis parvula</i>	<i>Leptosphaeria culmorum</i>	Connors (1967)
<i>Eleocharis pauciflora</i>	<i>Septoria punctoidea</i>	Connors (1967)
<i>Elodea canadensis</i>	<i>Alternaria tenuissima</i>	Bernhardt (1983)
	<i>Phythium afertile</i>	Bernhardt (1983)
<i>Hydrilla verticillata</i>	<i>Aspergillus awomori</i>	Pennington (1985)
	<i>Aspergillus</i> sp.	Freeman (1977)
	<i>Cephalosporium</i>	Charudattan (1973)
	<i>Chaetophoma</i> sp.	Charudattan (1973)
	<i>Curvularia</i> sp.	Varghese and Singh (1973)
	<i>Fusarium roseum</i>	Charudattan et al. (1984)
	<i>Fusarium solani</i>	Charudattan et al. (1978)
	<i>Gliocephalis</i>	Charudattan et al. (1978)
	<i>Humicola</i> sp. w.	Pennington (1985)
	<i>Trichoderma</i> sp.	Pennington (1985)
	<i>Oidiodendron</i> sp.	Charudattan (1973)
	<i>Penicillium</i> sp.	Freeman (1977)
	<i>Periconia</i> sp.	Charudattan (1973)
	<i>Phoma</i> sp.	Charudattan (1973)
	<i>Phytophthora parasitica</i>	Freeman (1977)
<i>Phythium</i> sp.	Varghese and Singh (1973)	
<i>Rhizoctonia solani</i>	Joyner and Freeman (1973)	
<i>Sclerotium rolfsi</i>	Freeman (1977)	
<i>Stigmina</i> sp.	Charudattan (1973)	

(Continued)

(Sheet 2 of 7)

Table 3 (Continued)

Plant Species	Fungus	Reference
<i>Hydrilla</i> <i>verticillata</i> (Continued)	<i>Trichoderma</i> sp.	Freeman (1977)
	<i>Verticillium</i> sp.	Varghese and Singh (1973)
<i>Myriophyllum</i> <i>spicatum</i>	<i>Acremonium curvulum</i>	Andrews et al. (1982)
	<i>Alternaria tenuissima</i>	Bernhardt (1983)
	<i>Cephalosporium</i> sp.	Bernhardt (1983)
	<i>Chaetomium funicolum</i>	Bernhardt (1983)
	<i>Cylindrocarpon</i> sp.	Bernhardt (1983)
	<i>Dactylella microaquatica</i>	Hayslip and Zettler (1973)
	<i>Epicoccum nigrum</i>	Bernhardt (1983)
	<i>Flagellospora curvula</i>	Bernhardt (1983)
	<i>Flagellospora stricta</i>	Hayslip and Zettler (1973)
	<i>Fusarium moniliforme</i>	Hayslip and Zettler (1973)
	<i>Fusarium nivale</i>	Bernhardt (1983)
	<i>Fusarium oxysporium</i>	Hayslip and Zettler (1973)
	<i>Fusarium roseum</i>	Hayslip and Zettler (1973)
	<i>Fusarium solani</i>	Hayslip and Zettler (1973)
	<i>Fusarium sporotrichoides</i>	Andrews and Hecht (1981)
	<i>Fusarium tricinctum</i>	Bernhardt (1983)
	<i>Gliocladium roseum</i>	Bernhardt (1983)
	<i>Jycoleptodiscus terrestris</i>	Gunner (1983)
	<i>Phoma</i> sp.	Bernhardt (1983)
	<i>Physoderma</i> sp.	Sparrow (1974)
	<i>Pythium afertile</i>	Bernhardt (1983)
	<i>Pythium carolinianum</i>	Bernhardt (1983)
	<i>Rhizoctonia solani</i>	Freeman et al. (1981)
	<i>Sclerotium hydrophyllum</i>	Hayslip and Zettler (1973)
	<i>Trichoderma</i> sp.	Bernhardt (1983)
	<i>Trichothecium</i> sp.	Andrews (1980)
	<i>Najas flexilis</i>	<i>Rhizoctonia solani</i>

(Continued)

(Sheet 3 of 7)

Table 3 (Continued)

Plant Species	Fungus	Reference
<i>Nuphar advena</i>	<i>Entyloma nymphaeae</i>	Index of Plant Diseases
	<i>Mycosphaerella pontederiae</i>	Index of Plant Diseases
	<i>Phyllosticta fatiscens</i>	Index of Plant Diseases
	<i>P. nymphaeacea</i>	Index of Plant Diseases
	<i>Sporobolomyces roseus</i>	Connors (1967)
<i>Nuphar luteum</i>	<i>Cercospora numphaeacea</i>	Charudattan et al. (1978)
	<i>Dichotomophothoropsis numphaerum</i>	Johnson and King (1976)
	<i>Pythium</i> sp.	Van der Velde (1978)
	<i>Sclerotium</i> sp.	Johnson et al. (1976)
<i>Nymphaea odorata</i>	<i>Alternaria</i> sp.	Horst (1979)
	<i>Cercospora exotica</i>	Horst (1979)
	<i>Cercospora nymphaeacea</i>	Index of Plant Diseases
	<i>Dichotomophothoropsis numphaearum</i>	Johnson and King (1976)
	<i>Entyloma nymphaeae</i>	Index of Plant Diseases
	<i>Mycosphaerella pontederiae</i>	↓
	<i>Ovularia numphaerum</i>	
	<i>Phyllosticta fatiscens</i>	
	<i>Pythium</i> sp.	
<i>Sclerotium</i> sp.	Johnson et al. (1976)	
<i>Nymphaea tuberosa</i>	<i>Dichotomophothoropsis numphaerum</i>	Johnson and King (1976)
	<i>Entyloma nymphaeae</i>	Index of Plant Diseases
	<i>Gleosporium numphaearum</i>	Connors (1967)
	<i>Pythium</i> sp.	Index of Plant Diseases
<i>Nymphoides peltata</i>	<i>Puccinia scirpi</i>	Munda and Garrasi (1978)
	<i>Septoria villarsiae</i>	Munda and Garrasi (1978)
<i>Nymphoides cordata</i>	<i>Burrillia decipiens</i>	Connors (1967)
	<i>Burrillia limnanthemii</i>	Connors (1967)
<i>Oenanthe sarmentosa</i>	<i>Nyssopsora echinata</i>	Connors (1967)
	<i>Septoria oenanthis</i>	Connors (1967)
<i>Phragmites communis</i>	<i>Claviceps purpurea</i>	↓
	<i>Graphyllum graminis</i>	
	<i>Hendersonia arundinacea</i>	
	<i>Mycosphaerella</i> sp.	

(Continued)

(Sheet 4 of 7)

Table 3 (Continued)

Plant Species	Fungus	Reference
<i>Phragmites communis</i> (Continued)	<i>Napicladium arundinaceum</i> <i>Neovossia iowensis</i> <i>Papularia arundinis</i> <i>Puccinia mogrusiana</i> <i>Puccinia phragmitis</i> <i>Pythium debaryanum</i> <i>Pythium ultimum</i> <i>Scolecotrichum graminis</i> <i>Selenophoma graminis</i> <i>Stagonospora</i> spp.	Index of Plant Diseases ↓
<i>Pistia stratiotes</i>	<i>Rhizoctonia solani</i>	Joyner and Freeman (1973)
<i>Pontederia lanceolata</i>	<i>Rhizoctonia solani</i> <i>Uromyces pontederiae</i>	Joyner and Freeman (1973) Charudattan (1981)
<i>Pontederia cordata</i> <i>Potamogeton crispus</i>	<i>Cercospora pontederiae</i> <i>Coniothyrium</i> sp. <i>Dactylella aquatica</i> <i>Geotrychium</i> sp. <i>Papulaspora aspera</i> <i>Pythium afertile</i> <i>Trichoderma</i> sp.	Conners (1967) Bernhardt (1983) Bernhardt (1983) Bernhardt (1983) Bernhardt and Duniway (1985) Bernhardt (1983) Bernhardt (1983)
<i>Potamogeton natans</i>	<i>Doassansia hydrophila</i>	Lowe (1969)
<i>Potamogeton nodosus</i>	<i>Curvularia</i> sp. <i>Doassansia martanoffiana</i> <i>Papulaspora aspera</i> <i>Pythium</i> sp.	Bernhardt (1983) Conners (1967) Bernhardt and Duniway (1985) Bernhardt (1983)
<i>Potamogeton pectinatus</i>	<i>Anquillospora crassa</i> <i>Cephalosporium</i> sp. <i>Curvularia</i> sp. <i>Diplosporium</i> sp. <i>Flagellospora curvula</i> <i>Fusarium moniliforme</i>	Bernhardt (1983) Bernhardt (1983) Lumsden et al. (1963) Bernhardt (1983) Bernhardt (1983) Bernhardt (1983)

(Continued)

(Sheet 5 of 7)

Table 3 (Continued)

Plant Species	Fungus	Reference
<i>Potamogeton pectinatus</i>	<i>Hyaloflorae</i> sp.	Lumsden et al. (1963)
	<i>Nigrospora oryzae</i>	Bernhardt (1983)
	<i>Phoma</i> sp.	Lumsden et al. (1963)
	<i>Pullaria pullulans</i>	Lumsden et al. (1963)
	<i>Pythium afertile</i>	Bernhardt (1983)
	<i>Pythium carolinianum</i>	Bernhardt (1983)
	<i>Pythium</i> sp.	Lumsden et al. (1963)
	<i>Rhizoctonia solani</i>	Bourn and Jenkins (1928)
<i>Potamogeton perfoliatus</i>	<i>Cephalosporium</i> sp.	Motta (1978)
	<i>Papulaspora</i> sp.	Motta (1978)
	<i>Pythium</i> sp.	Motta (1978)
	<i>Rhizoctonia solani</i>	Bourn and Jenkins (1928)
	<i>Trichoderma viride</i>	Motta (1978)
<i>Potamogeton</i> sp.	<i>Doassansia martianoffiana</i>	Index of Plant Diseases
	<i>D. occulta</i>	
	<i>Lagenidium muenschari</i>	
	<i>Ramularia aquatilis</i>	
	<i>Rhizoctonia solani</i>	
<i>Ruppia cirrhosa</i>	<i>Tetramyxa parasitica</i>	Verhoeven (1975)
<i>Ruppia maritima</i>	<i>Cephalosporium</i> sp.	Motta (1978)
	<i>Papulospora</i> sp.	Motta (1978)
	<i>Pythium</i> sp.	Motta (1978)
	<i>Rhizoctonia solani</i>	Bourn and Jenkins (1928)
	<i>Trichoderma viride</i>	Motta (1978)
<i>Salvinia auriculata</i>	<i>Alternaria</i> sp.	Loveless (1969)
	<i>Spicariopsis</i> sp.	Loveless (1969)
<i>Salvinia rotundifolia</i>	<i>Rhizoctonia solani</i>	Joyner and Freeman (1973)
<i>Sagittaria</i> sp.	<i>Cercospora alismatis</i>	Horst (1979)
	<i>Cercospora sagittariae</i>	Connors (1967)
	<i>Didymaria alismatis</i>	↓
	<i>Gleosporium confluens</i>	
	<i>Marssonina</i> sp.	
	<i>Burrillia pustulate</i>	
	<i>Doassansia deformans</i>	
	<i>Doassansia furva</i>	
<i>Doassansia intermedia</i>		
	(Continued)	

Table 3 (Concluded)

Plant Species	Fungus	Reference
<i>Sagittaria</i> sp. (Continued)	<i>Doassansia obscura</i> <i>Doassansia opaca</i> <i>Doassansia sagittariae</i> <i>Fusarium acuminatum</i> <i>Rhynchosporium alismatis</i>	Connors (1967) ↓
<i>Scirpus articulatus</i>	<i>Physoderma thilumalacharii</i>	Pavgi and Singh (1980)
<i>Scirpus</i> sp.	<i>Heterosporium</i>	Lowe (1969)
<i>Spartina</i> <i>alterniflora</i>	<i>Buergenerula spartinae</i> <i>Sphaerulina pedicellata</i> <i>Uromyces acuminatus</i>	Torzilli and Andrykovitch (1980) Gessner et al. (1972) Connors (1967)
<i>Spirodela polyrhiza</i>	<i>Tracya lemnae</i>	Connors (1967)
<i>Typha latifolia</i>	<i>Cladosporium</i> spp. <i>Cryptomela typhae</i> <i>Hendersonia typhae</i> <i>Heterosporium maculatum</i> <i>Hymenopsis hydrophila</i> <i>Leptosphaeria eustoma</i> <i>Leptosphaeria typharum</i> <i>Lophodermium typhinum</i> <i>Mycosphaerella typhae</i> <i>Ophiobolus</i> sp. <i>Phyllosticta typhina</i> <i>Pleospora typhae</i> <i>Pythiogeton autossytum</i> <i>Pythium helicoides</i> <i>Scolecotrichium typhae</i> <i>Stagnospora typhiodearum</i>	Index of Plant Diseases Index of Plant Diseases Index of Plant Diseases Lowe (1969) Index of Plant Diseases Horst (1979) Index of Plant Diseases ↓
<i>Vallisneria spiralis</i>	<i>Rhizoctonia solani</i>	Bourn and Jenkins (1928)
<i>Zostera marina</i>	<i>Acremonium</i> sp. <i>Cladosporium</i> sp. <i>Labyrinthula</i> sp. <i>Lulworthia</i> sp. <i>Ophiobolus halimus</i>	Newel (1981) Newel (1981) Renn (1936) Newel (1981) Tutin (1938)

Table 4

List of Selected Plant-Parasitic Nematodes
Associated with Aquatic Plants

Plant Species	Nematode	Reference	
<i>Alisma plantago aquatica</i>	<i>Heterodera aquatica</i>	Gerber et al. (1986b)	
<i>Altermanthera phyloxeroïdes</i>	<i>Cacopaurus</i>	Esser et al. (1985) ↓ Plakidas (1936) Index of Plant Diseases Esser et al. (1985) Esser et al. (1985) Esser et al. (1985) Esser et al. (1985)	
	<i>Criconemoides</i>		
	<i>Dolichodorus</i>		
	<i>Helicotylenchus crenacauda</i>		
	<i>Helicotylenchus erythrinae</i>		
	<i>Heterodera marioni</i>		
	<i>Meloidogyne</i> sp.		
	<i>Pratylenchus</i>		
	<i>Trichodorus proximus</i>		
	<i>Tylenchus</i>		
<i>Xiphinema</i>			
<i>Ceratophyllum demersum</i>	<i>Aphelenchoides fragariae</i>	Smart and Esser (1968) Gerber et al. (1986a) ↓	
	<i>Aphelenchoides</i> sp.		
	<i>Criconemoides</i> sp.		
	<i>Hirschmanniella caudacrena</i>		
	<i>Hirschmanniella</i> sp.		
	<i>Hopolaimus tylenchiformis</i>		
	<i>Pratylenchus</i> sp.		
	<i>Trichodorus</i> sp.		
<i>Tylenchus</i> sp.			
<i>Eichhornia crassipes</i>	<i>Xiphidorus amazonensis</i>	Gerber et al. (1986b)	
<i>Eleocharis acicularis</i>	<i>Helicotylenchus</i> sp.	↓	
	<i>Heterodera canadensis</i>		
	<i>Meloidogyne sewelli</i>		
<i>Eleocharis palustris</i>	<i>Meloidogyne</i> sp.		
<i>Eleocharis</i> sp.	<i>Doplichodorus similis</i>		
<i>Elodea canadensis</i>	<i>Anguillula</i> sp.		
	<i>Aphelenchus</i> sp.		
	<i>Hirschmanniella oryzae</i>		
<i>Hydrilla verticillata</i>	<i>Aphelenchoides fragariae</i>		Gerber et al. (1986a) ↓
	<i>Aphelenchoides</i> sp.		
	<i>Criconemoides</i> sp.		

(Continued)

(Sheet 1 of 4)

Table 4 (Continued)

Plant Species	Nematode	Reference
<i>Hydrilla</i> <i>verticillata</i> (Continued)	<i>Dolichodorus</i> sp. <i>Helicotylenchus</i> sp. <i>Hirschmanniella caudacrena</i> <i>Hirschmanniella gracilis</i> <i>Hirschmanniella oryzae</i> <i>Hirschmanniella</i> sp. <i>Trichodorus</i> sp. <i>Thylenchorhynchus irregularis</i> <i>Tylenchus</i> sp. <i>Xiphinema americanum</i>	Gerber et al. (1986a) ↓
<i>Lemna</i> sp.	<i>Aphelenchoides fragariae</i>	Smart and Esser (1968)
<i>Limnium spongia</i>	<i>Aphelenchoides fragariae</i>	Esser et al. (1985)
<i>Lythrum salicaria</i>	<i>Meloidogyne</i> sp.	Gerber et al. (1986b)
<i>Myriophyllum</i> <i>aquaticum</i>	<i>Aphelenchoides</i> sp. <i>Hirschmanniella</i> sp.	Esser et al. (1985) Esser et al. (1985)
<i>Myriophyllum</i> <i>spicatum</i>	<i>Aphelenchoides fragariae</i> <i>Hirschmanniella gracilis</i>	Gerber et al. (1986b) Gerber et al. (1986b)
<i>Myriophyllum</i> <i>verticillatum</i>	<i>Ditylenchus dipsaci</i>	Gaevskaya (1969)
<i>Najas flexilis</i>	<i>Aphelenchoides fragariae</i>	Gerber et al. (1986b)
<i>Najas</i> spp.	<i>Aphelenchoides fragariae</i> <i>Hirschmanniella caudacrena</i>	Smart and Esser (1968) Esser et al. (1985)
<i>Nuphar luteum</i>	<i>Chronogaster</i> <i>Chrysonemoides</i> <i>Dorylaimoides</i> <i>Plectus</i> <i>Tobrilus</i> <i>Tripyla</i>	Prejs (1977) ↓
<i>Nymphaea</i> sp.	<i>Criconemoides</i> <i>Dolichodorus heterocephallus</i> <i>Helicotylenchus</i> <i>Meloidogyne</i> <i>Pratylenchus</i> <i>Rotylenchus</i> <i>Trichodorus</i>	Esser et al. (1985) ↓

(Continued)

(Sheet 2 of 4)

Table 4 (Continued)

Plant Species	Nematode	Reference
<i>Oenanthe</i> sp.	<i>Meloidogyne marioni</i>	Gaevskaya (1969)
<i>Phragmites communis</i>	<i>Anguina graminis</i>	Gaevskaya (1969)
	<i>Ditylenchus radicicola</i>	Gerber et al. (1986b)
	<i>Dolichodorus</i> sp.	Gerber et al. (1986b)
	<i>Helicotylenchus erythrinae</i>	Gerber et al. (1986b)
	<i>Hirschmanniella gracilis</i>	Gerber et al. (1986b)
<i>Pistia stratiotes</i>	<i>Hirschmanniella caudacrena</i>	Esser et al. (1985)
<i>Pontederia lanceolata</i>	<i>Dolichodorus</i>	Esser et al. (1985)
	<i>Tylenchorhynchus</i>	Esser et al. (1985)
<i>Potamogeton friesii</i>	<i>Ditylenchus dipsaci</i>	Gaevskaya (1969)
<i>Potamogeton lucens</i>	<i>Calolaimus</i>	Prejs (1986)
	<i>Chromadorita</i>	Prejs (1977)
	<i>Chronogaster</i>	Prejs (1977)
	<i>Chrysonemoides</i>	Prejs (1977)
	<i>Cryptonchus</i>	Prejs (1986)
	<i>Dorylaimoides</i>	Prejs (1977)
	<i>Dorylaimus</i>	
	<i>Hirschmanniella</i>	
	<i>Panagrolaimus</i>	
	<i>Plectus</i>	
	<i>Tobrilus</i>	
	<i>Tripyla</i>	
<i>Potamogeton pectinatus</i>	<i>Hirschmanniella</i>	Prejs (1986)
	<i>Panagrolaimus</i>	
	<i>Plectus</i>	
<i>Potamogeton perfoliatus</i>	<i>Calolaimus</i>	
	<i>Chronogaster</i>	
	<i>Chrysonemoides</i>	
	<i>Hirschmanniella</i>	
	<i>Cryptonchus</i>	
	<i>Panagrolaimus</i>	
<i>Plectus</i>		
<i>Potamogeton</i> sp.	<i>Aphelenchoides fragariae</i>	Smart and Esser (1968)
<i>Ruppia maritima</i>	<i>Aphelenchoides</i>	Esser et al. (1985)
	<i>Helicotylenchus</i>	
	<i>Hirschmanniella gracilis</i>	

(Continued)

(Sheet 3 of 4)

Table 4 (Concluded)

Plant Species	Nematode	Reference
<i>Ruppia maritima</i> (Continued)	<i>Hirschmanniella oryzae</i> <i>Tylenchus</i>	Esser et al. (1985) Esser et al. (1985)
<i>Sagittaria</i> <i>sagittifolia</i>	<i>Chrysonemoides</i> <i>Dorylaimus</i> <i>Plectus</i>	Prejs (1977) Prejs (1977) Prejs (1977)
<i>Sagittaria</i> sp.	<i>Aphelenchoides</i> <i>Dolichodorus</i> <i>Helicotylenchus</i> <i>Hirschmanniella caudacrena</i> <i>Scutellonema</i> <i>Tylenchus</i> <i>Xiphinema</i>	Esser et al. (1985) ↓
<i>Sirpus americanus</i>	<i>Hirschmanniella anchoryzae</i> <i>Hirschmanniella gracilis</i>	Gerber et al. (1986b) Gerber et al. (1986b)
<i>Spartina</i> <i>alterniflora</i>	<i>Oncholaimus</i> <i>Meloidogyne spartinae</i>	Gessner et al. (1972) Gerber et al. (1986b)
<i>Typha</i> sp.	<i>Criconemoides mutabile</i> <i>Dolichodorus</i> spp. <i>Helicotylenchus pseudorobustus</i> <i>Hemicriconemoides</i> sp. <i>Hemicycliophora</i> sp. <i>Heterodera</i> sp. <i>Hirschmanniella belli</i> <i>Hirschmanniella spinicaudata</i> <i>Hirschmanniella</i> sp. <i>Medloidyne marioni</i> <i>Pratylenchus neglectus</i> <i>Pratylenchus thornei</i> <i>Trichodorus</i> sp. <i>Thylenchorhynchus clarus</i> <i>Tylenchulus semipenetrans</i>	Siddiqui et al. (1973) ↓ Gaevs kaya (1969) Siddiqui et al. (1973) ↓
<i>Vallisneria</i> <i>americana</i>	<i>Hirschmanniella caudacrena</i> <i>Tylenchorhynchus irregularis</i> <i>Tylenchus</i> sp. <i>Xiphinema americanum</i>	Esser et al. (1985) Gerber et al. (1986b) Gerber et al. (1986b) Gerber et al. (1986b)

Table 5
Senescence-Inducing Factors

<u>Inducing Factor</u>	<u>Manipulation in the Field*</u>	
	<u>Feasible</u>	<u>Infeasible</u>
Low light	++	
Nutrient availability		+
Photoperiod	++	
Temperature		+
Dense periphyton	+	
External growth regulators (ABA)	++	
Sublethal herbicide doses	+++	
Herbivores	++	

* + = low, ++ = high, +++ = higher feasibility.

APPENDIX A: GLOSSARY*

- Commensalism--The living together of two or more organisms with benefit usually to one and without injury to either.
- Decomposition--Degradation (often microbial) of dead and organic matter.
- Disease--Injurious alteration of one or more ordered processes of energy utilization in a living system caused by the continued irritation of a primary causal factor or factors.
- Epiphytic--Being attached to the surface of an organism (plant).
- Epiphyton--The assemblage of (micro)organisms that grow attached to aquatic macrophytes.
- Epiphytotic--Relating to a widespread and destructive disease of plants (epidemic).
- Host (suscept)--The plant that is being acted on by a pathogen.
- Host range of pathogen--The kinds of plants a specific pathogen can infect and cause disease or set up an infection process.
- Infection--The process in which a parasite or biotic pathogen establishes contact with the host tissues.
- Infectious diseases--Those caused by living organisms (e.g., fungi or bacteria) in contrast to noninfectious disease caused by nonliving pathogens (air pollutants).
- Inoculum--The infective unit or propagule (e.g., the spore) that is carried to the host plant and initiates infection.
- Parasite--An organism that obtains food from another living organism (host).
- Pathogenesis--Process of disease development in the host.
- Pathogenicity--The ability of a pathogen to produce disease in a given host.
- Pathogens--Factors or agents that cause plant disease; may be living (biotic) or nonliving (abiotic).
- Periphyton--Assemblage of organisms growing upon free surfaces of submerged objects in water (corresponds to the German term "Aufwuchs," which is often used in English terminology).
- Saprophyte--An organism that can obtain nutrients from dead organic material.
- Senescence--The final phase in development of an organism leading to cellular breakdown and death.
- Symptoms--Outward manifestations of disease.
- Vector--An agent (biotic or abiotic) able to transmit a pathogen.
- Virulence--The degree of pathogenicity.

* Based on definitions given by Hanson (1957), Holcomb (1982), Lucas et al. (1985), and Sladeckova (1962).